

THE COMPUTATION OF GALOIS GROUPS

Leonard Soicher

A Thesis
in
The Department
of
Computer Science

Presented in Partial Fulfillment of the Requirements
for the degree of Master of Computer Science at
Concordia University
Montreal, Quebec, Canada

April, 1981

© Leonard Soicher, 1981

ABSTRACT

THE COMPUTATION OF GALOIS GROUPS

Leonard Soicher

We discuss methods of computing invariants of the conjugacy class of the Galois group of a separable polynomial $f(x)$ over K , $n = \deg(f) > 0$. The aim is to determine the class of $\text{Gal}(f/K)$ in S_n . We concentrate on the case $K = Q$ and $f(x)$ is irreducible over K .

The main tool discussed is the resolvent polynomial. For F in $K[x_1, \dots, x_n]$, the factorization of a resolvent polynomial is used to determine the orbit length partition of $\{F(x_{1P}, \dots, x_{nP}) : P \in S_n\}$ under the action of $\text{Gal}(f/K)$.

An important class of resolvent polynomials considered are the "linear" resolvent polynomials, where $F = e_1x_1 + \dots + e_rx_r$, e_i in K and $0 < r \leq n$. The use of linear resolvents in determining $\text{Gal}(f/K)$ is discussed. A new, practical, exact method of computing linear resolvents is described, as well as the computer implementation of this method over the integers.

For every transitive permutation group G of degree up to 7, we have computed a polynomial $f(x)$ such that $\text{Gal}(f/Q) = G$. We also list many new examples of polynomials with $\text{PSL}(3,2)$ as Galois group over Q .

ACKNOWLEDGEMENTS

I would like to thank my teacher and supervisor, Prof. J. McKay, for his superb guidance and support throughout this work.

I also thank the following people who made their results and/or computer programs available to me: G. Butler, D. Ford, G. Kolesova, H. Kisilevsky, E. Regener and R. Rohlicek.

Also, to my friends and family I say thank you for your encouragement and support.

This research was funded by the Natural Sciences and Engineering Research Council of Canada and by the Government of Quebec.

TABLE OF CONTENTS

ABSTRACT

ACKNOWLEDGEMENTS

TABLE OF CONTENTS

LIST OF TABLES

NOTATION

CHAPTER 1. INTRODUCTION.....1

 1.1 Basic Definitions and Results2

 1.1.1 Group Actions2

 1.1.2 The Galois Group of a Polynomial4

 1.1.3 The Fundamental Theorem of Galois Theory6

 1.1.4 The Fundamental Theorem on Symmetric Polynomials7

 1.1.5 The Resolvent Polynomial8

 1.1.6 The Discriminant9

 1.1.7 The Specialization to \mathbb{Q} 10

 1.2 Content and Contribution of this Thesis10

CHAPTER 2. METHODS OF DETERMINING GALOIS GROUPS.....13

 2.1 Determining the Galois Group in Finitely Many Steps13

 2.2 The Determination of Cycle Types in $\text{Gal}(f/\mathbb{Q})$ 15

 2.3 The Resolvent Polynomial18

 2.3.1 Theoretical Development18

 2.3.2 Construction and Factorization of Resolvents22

2.3.3	Functions Belonging to Groups	24
2.3.3.1	The Alternating Function	24
2.3.4	The Method of Stauduhar	25
2.3.5	The Use of Linear Resolvent Polynomials	27
2.3.5.1	Action on Sets and Sequences	28
2.3.5.2	Differentiating all Transitive Groups of Degree up to 7	30
CHAPTER 3. LINEAR RESOLVENT POLYNOMIAL CONSTRUCTION....		33
3.1	Restrictions on the Field	33
3.2	Polynomial Operations	34
3.2.1	The Greatest Common Divisor	34
3.2.2	The Resultant	35
3.2.3	The Formal Derivative and its Zeros	37
3.2.4	"Multiply Zeros"	38
3.2.5	"Sum Zeros"	38
3.2.6	"Polynomial Root"	39
3.3	Multiset Operations	40
3.4	Constructive Proof	41
3.5	Algorithm LINRESOLV	42
3.6	Remarks	45
CHAPTER 4. IMPLEMENTATION AND EXAMPLES.....		47
4.1	A Modular Approach to Computing Resolvents	47
4.1.1	Bounding the Zeros of $f(x)$	48
4.2	The Implementation	49
4.2.1	LINRESOLV over $K = \mathbb{Z}_p$	49
4.3	Examples	51

REFERENCES	54
APPENDIX 1. TABLES OF TRANSITIVE GROUPS OF DEGREE UP TO 8 (SUPPLIED BY G. BUTLER).	56
APPENDIX 2. POLYNOMIALS WITH GIVEN TRANSITIVE GALOIS GROUPS OVER Q OF DEGREE UP TO 7	85
APPENDIX 3. POLYNOMIALS WITH $PSL(3,2)$ AS GALOIS GROUP OVER Q	90

LIST OF TABLES

APPENDIX 1

3A	Groups of Degree 3	59
3B	Group Generators	59
3C	Cycle Type Distribution	59
3D	Orbit Length Partitions of Sets and Sequences under G	60
4A	Groups of Degree 4	61
4B	Group Generators	61
4C	Cycle Type Distribution	61
4D	Orbit Length Partitions of Sets and Sequences under G	62
5A	Groups of Degree 5	63
5B	Group Generators	63
5C	Cycle Type Distribution	63
5D	Orbit Length Partitions of Sets and Sequences under G	64
6A	Groups of Degree 6	65
6B	Group Generators	66
6C	Cycle Type Distribution	67
6D	Orbit Length Partitions of Sets and Sequences under G	68
7A	Groups of Degree 7	69
7B	Group Generators	69
7C	Cycle Type Distribution	70
7D	Orbit Length Partitions of Sets and Sequences under G	71
8A	Groups of Degree 8	72
8B	Group Generators	74

8C	Cycle Type Distribution	76
8D	Orbit Length Partitions of Sets and Sequences under G	82

APPENDIX 2

A2.1	Polynomials $f(x)$ such that	88
	$\text{Gal}(f/Q) = G$	

APPENDIX 3

A3.1	Polynomials $f(x)$ such that	91
	$\text{Gal}(f/Q) = \text{PSL}(3, 2)$	

NOTATION

tS	the image of t under the mapping S ,
$ C $	the cardinal of the set C ,
K	a field,
$\text{char}(K)$	the characteristic of K ,
$K(v_1, \dots, v_n)$	the field extension of K obtained by adjoining v_1, \dots, v_n to K ,
$R[x_1, \dots, x_n]$	the polynomials in the indeterminates x_1, \dots, x_n with coefficients in R ,
$f(x)$	a polynomial in $K[x]$,
$\deg(f)$	the degree of $f(x)$,
$\text{disc}(f)$	the discriminant of $f(x)$,
$f'(x)$	the formal derivative of $f(x)$,
$G(N/K)$	the Galois group of the normal extension N over K ,
$\text{Gal}(f/K)$	the Galois group of $f(x)$ over K ,
Q	the field of rational numbers,
Z	the ring of rational integers,
p	a positive prime,
Z_p	the field of integers modulo p ,
$i \bmod p$	the image of i under the natural homomorphism from Z onto Z_p ,
$F \bmod p$	F with its coefficients replaced by their images mod p (F a (multivariate) polynomial with coefficients in Z),

$f \bmod g$	the remainder upon division of $f(x)$ by $g(x)$,
$\text{res}(f,g)$	the resultant of $f(x)$ and $g(x)$,
G	a group,
S_n	the symmetric group on $\{1, \dots, n\}$,
A_n	the alternating group on $\{1, \dots, n\}$,
$H < G$	H is a subgroup of G ,
$*$	a group action,
$\text{rep}(G, \underline{C}, *)$	the representation of G into S_n defined by the action of G by $*$ on the ordered set \underline{C} ,
$\text{im}(\text{rep}(G, \underline{C}, *))$	the image of G under the preceding representation,
$\text{stab}_G(c)$	the stabilizer in G of c ,
$[e_1, \dots, e_r]$	the multiset of elements e_1, \dots, e_r ,
$\text{mult}(e, M)$	the multiplicity of the element e in the multiset M ,
$\text{mult}(v, f)$	the multiplicity of v as a zero of $f(x)$,
$\text{gcd}(a, b)$	the greatest common divisor of a and b ,
$a b$	a divides b ,
$a b$	a divides b and $\text{gcd}(a, b/a) = 1$,
$a <- \text{expression}$	the value of a is replaced by the value of the expression.

CHAPTER 1

INTRODUCTION

Galois theory gives an elegant answer to the question of whether a polynomial equation, $f(x) = 0$, over a suitable field K (e.g. the rationals) is solvable by radicals.

"Solvable by radicals" means that the zeros of $f(x)$ can be expressed as finite expressions in the coefficients of $f(x)$, where the only permitted operations are the field operations and the extraction of roots. In Galois theory, to each polynomial $f(x)$ over K , there is an associated group G called the Galois group of $f(x)$ over K . The structure of this group describes the structure of the smallest field extension of K containing all the zeros of $f(x)$, and the equation $f(x) = 0$ is solvable by radicals if and only if G is a solvable group [VDW, p.173].

In this thesis we study the problem of computing the Galois group of a given polynomial $f(x)$, with distinct zeros, over a field K . We are especially interested in the case $K = \mathbb{Q}$, the field of rational numbers, and when $f(x)$ is irreducible over K . The thesis is intended as a contribution to the domain of symbolic and algebraic computation.

We assume the reader is familiar with basic algebra including group theory, field extension theory and Galois theory. References for this algebra are [BIR, VDW].

1.1. BASIC DEFINITIONS AND RESULTS

We define our terms and state several useful basic results.

1.1.1 GROUP ACTIONS

We define the action of a group on a set. This is fundamental as we will be concerned with determining the action of the Galois group on various sets.

DEFINITION 1.1. Let C be a set and G be a group. We say that G acts on C (by $*$), if for each pair (c, s) where c in C and s in G , there is defined an element $c*s$ in C such that for all c in C and s, t in G the following axioms hold:

- (1) $c*1_G = c$, where 1_G is the identity element of G , and
- (2) $(c*s)*t = c*(st)$.

Let G be a group, C a set, and suppose G acts on C by $*$. Let c be in C .

The orbit containing c (under G) is defined by

$$c*G = \{c*s : s \text{ in } G\}.$$

$|c*G|$ is called the orbit length. The set of orbits of C

under G ,

$$\{c^*G : c \text{ in } C\},$$

partitions C . This partition of C induces a partition of $|C|$, called the orbit length partition of C under G . This partition of $|C|$ consists of the lengths of the distinct orbits of C under G .

The stabilizer of c in G is defined by

$$\text{stab}_G(c) = \{S \text{ in } G : c^*S = c\}.$$

Let S, T in G , c, d in C and $H = \text{stab}_G(c)$. It is straightforward to show that the following facts are true (see [NEU]):

- (1) $c^*S = c^*T$ if and only if $HS = HT$; that is, iff S and T are in the same right coset of $\text{stab}_G(c)$ in G .
- (2) $\text{stab}_G(c^*S) = S^{-1}HS$.
- (3) Suppose that $|C| = n < \infty$, and let an ordering of the elements of C be $\underline{C} = (c_1, \dots, c_n)$. Then there is a natural permutation representation (homomorphism):

$$\text{rep}(G, \underline{C}, *) : G \rightarrow S_n,$$

where $S \rightarrow \overline{S}$ under this representation, and \overline{S} is defined by:

$$i\overline{S} = j \text{ if and only if } c_i^*S = c_j,$$

for all S in G and i in $\{1, \dots, n\}$. The kernel of $\text{rep}(G, \underline{C}, *)$ is

$$\bigcap_{i=1}^n \text{stab}_G(c_i).$$

The subgroup of S_n which is the image of $\text{rep}(G, \underline{C}, *)$ is denoted by $\text{im}(\text{rep}(G, \underline{C}, *))$.

Let $H = \text{im}(\text{rep}(G, \underline{C}, *))$, and let P be a permutation in S_n . Consider a new ordering of the elements of C :

$$\underline{C}' = (c'_1, \dots, c'_n) = (c_{1P}, \dots, c_{nP}).$$

Then $\text{im}(\text{rep}(G, \underline{C}', *)) (= PHP^{-1})$.

1.1.2 THE GALOIS GROUP OF A POLYNOMIAL

Let $f(x) = \sum_{i=0}^n a_i x^i$ be a polynomial in $K[x]$, $a_n \neq 0$, $n = \deg(f) > 0$.

DEFINITION 1.2. We say that a field extension N of K is a splitting field of $f(x)$ over K if:

- (1) $f(x)$ can be factored into linear factors, $f(x) = a_n(x-v_1)\dots(x-v_n)$, in $N[x]$, and
- (2) N is generated over K by v_1, \dots, v_n , that is, $N = K(v_1, \dots, v_n)$.

We call v_1, \dots, v_n the zeros of $f(x)$, and we may assume that $f(x)$ is monic ($a_n = 1$).

From field-extension theory we know that for the given field K , and $f(x)$ in $K[x]$, we can always construct a splitting field of $f(x)$ over K and this splitting field is unique up to field isomorphism. Thus we may speak of the

splitting field of $f(x)$ over K .

DEFINITION 1.3. Let L be a field. An automorphism of L is a 1-to-1 mapping, S , of L onto L such that for all elements a, b in L , $(ab)S = (aS)(bS)$ and $(a+b)S = (aS)+(bS)$.

DEFINITION 1.4. Let N be the splitting field of $f(x)$ over K . The Galois group of N over K , denoted by $G(N/K)$, is the group of all the automorphisms of N which fix each element in K .

Let N be the splitting field of $f(x)$ over K and let $G = G(N/K)$. We call $f(x)$ separable if its zeros in N are distinct. Many of the results of Galois theory apply only to the splitting fields of separable polynomials (the so-called normal and separable extensions: if N is the splitting field of separable $f(x)$ over K , then each element w in N is a zero of a unique separable, monic, irreducible polynomial over K .) We now assume that $f(x)$ is a separable polynomial over K .

Let an ordering of the (distinct) zeros of $f(x)$ be $\underline{v} = (v_1, \dots, v_n)$. G sends a zero of $f(x)$ to a zero of $f(x)$ (see Lemma 2.6) and thus G acts on the set $V = \{v_1, \dots, v_n\}$ by $*$, where the action is defined by $v_i * S = v_i S$ for every S in G and i in $\{1, \dots, n\}$. Thus there is the natural representation

$$\text{rep}(G(N/K), \underline{v}, *) : G(N/K) \rightarrow S_n$$

as described in Section 1.1.1. This representation is

faithful since if an element T is in the kernel of $\text{rep}(G(N/K), \underline{v}, *)$, then T must fix each of the v_i as well as the elements of K . Since V generates N over K , T must be the identity element.

DEFINITION 1.5. The Galois group of $f(x)$ over K , $\text{Gal}(f/K)$, with respect to the ordering $\underline{v} = (v_1, \dots, v_n)$ of the zeros of $f(x)$, is defined to be $\text{im}(\text{rep}(G(N/K), \underline{v}, *))$.

If we do not fix an ordering of the zeros of $f(x)$, then $\text{Gal}(f/K)$ can be determined at best to within conjugacy in S_n . This is stronger than to within isomorphism and in this thesis we are usually not concerned with the problem of ordering the zeros of $f(x)$. If we do not specify an ordering of the zeros of $f(x)$ and we state that $\text{Gal}(f/K) = G$, we mean that for some ordering of the zeros of $f(x)$, $\text{Gal}(f/K) = G$ with respect to that ordering.

1.1.3. THE FUNDAMENTAL THEOREM OF GALOIS THEORY

For later reference, we state the Fundamental Theorem of Galois Theory (for a detailed discussion see [BIR]).

THEOREM 1.6. Let $G = G(N/K)$ be the Galois group of the splitting field N of a separable polynomial $f(x)$ over K . There is a 1-to-1 correspondence between the subgroups H of G and the subfields L of N which contain K . Given L , the corresponding subgroup H is the group of all the automorphisms in G which fix every element in L . Given H ,

the corresponding subfield L consists of all the elements of N left fixed by every automorphism in H . For each L , the corresponding subgroup H is the Galois group of N over L , and $|H|$ is the degree of N over L .

In particular, if an element b in N is left fixed by all automorphisms in $G(N/K)$, then b belongs to the base field K , the subfield of N corresponding to $G(N/K)$.

1.1.4 THE FUNDAMENTAL THEOREM ON SYMMETRIC POLYNOMIALS

We state the Fundamental Theorem on Symmetric Polynomials.

THEOREM 1.8. ([VDW, p.81]) Let R be a commutative ring with identity and let A in $R[x_1, \dots, x_n]$ be a symmetric polynomial (that is, $A(x_1, \dots, x_n) = A(x_{1p}, \dots, x_{nP})$ for every p in S_n). One can construct a unique polynomial B in $R[x_1, \dots, x_n]$ such that $A = B(s_1, \dots, s_n)$, where s_i is the i -th elementary symmetric polynomial (that is, $s_i = \sum_{j_1 < j_2 < \dots < j_i} x_{j_1} \dots x_{j_i}$, where the sum is taken over all $1 \leq j_1 < \dots < j_i \leq n$).

If monic $f(x) = \sum_{i=0}^n a_i x^{n-i}$ has zeros v_1, \dots, v_n , then $a_i = (-1)^i s_i(v_1, \dots, v_n)$, for $i=1, \dots, n$. Thus if R is a commutative ring with identity, then any symmetric polynomial over R in the zeros of $f(x)$ can be expressed as a polynomial over R in the coefficients of $f(x)$.

1.1.5 THE RESOLVENT POLYNOMIAL

Let $F = F(x_1, \dots, x_n)$ be a polynomial in $K[x_1, \dots, x_n]$ and let P be a permutation of $\{1, \dots, n\}$. We define

$$F^*P = F(x_{1P}, \dots, x_{nP}).$$

We call F^*P a conjugate function of F . In this way any permutation group on $\{1, \dots, n\}$ acts on $K[x_1, \dots, x_n]$ as a group of ring automorphisms.

DEFINITION 1.9. Let F be in $K[x_1, \dots, x_n]$, $f(x)$ in $K[x]$, and $n = \deg(f) > 0$. Let the zeros of $f(x)$ be v_1, \dots, v_n . The resolvent polynomial associated with F and $f(x)$, $R(F, f)$, is defined by:

$$R(F, f) = \prod_{i=1}^k (x - F_i(v_1, \dots, v_n)),$$

where $\{F_1, \dots, F_k\} = F^*S_n$ (F_i distinct functions).

We may take $F_i = F^*P_i$, ($i=1, \dots, k$), where $\{P_1, \dots, P_k\}$ is a set of right coset representatives of $\text{stab}_{S_n}(F)$ in S_n (see Section 1.1.1).

The coefficients of $R(F, f)$ are symmetric polynomials over K in v_1, \dots, v_n and hence by the Fundamental Theorem on Symmetric Polynomials, the coefficients of $R(F, f)$ can be expressed as polynomials over K in the coefficients of $f(x)$. We also note that $R(F, f)$ is independent of the ordering of the zeros of $f(x)$.

An important resolvent polynomial we consider in this thesis is what we call the linear resolvent polynomial.

DEFINITION 1.10. Let $f(x)$ be in $K[x]$, $n = \deg(f) > 0$, and let e_1, \dots, e_r be in K , $0 < r \leq n$. Let the multiset $M = [e_1, \dots, e_r]$. The linear resolvent polynomial associated with M and $f(x)$, $LR(M, f)$, is defined to be the resolvent polynomial associated with F and $f(x)$, when $F = F(x_1, \dots, x_n) = e_1x_1 + \dots + e_rx_r$.

1.1.6 THE DISCRIMINANT

An important symmetric function of the zeros of a polynomial $f(x)$ is the discriminant of $f(x)$.

DEFINITION 1.11. Let $f(x)$ be in $K[x]$, $n = \deg(f) > 1$, and let the zeros of $f(x)$ be v_1, \dots, v_n . The discriminant of $f(x)$, $\text{disc}(f)$, is defined by

$$\text{disc}(f) = \prod_{i < j} (v_i - v_j)^2.$$

We note that $\text{disc}(f) = 0$ if and only if the zeros of $f(x)$ are not distinct.

The discriminant of monic $f(x)$ can be computed efficiently using the relationship (see [CHI, p.283-286]):

$$(1.1) \quad \text{disc}(f) = (-1)^{n(n-1)/2} \text{res}(f, f')$$

where $\text{res}(f, f')$ is the resultant of $f(x)$ and its formal

derivative $f'(x)$. The resultant and formal derivative are discussed in Section 3.2.

1.1.7 THE SPECIALIZATION TO Q

Let monic separable $f(x)$ be in $\mathbb{Q}[x]$, $n = \deg(f) > 0$. We take the splitting field of $f(x)$ over \mathbb{Q} to be a subfield of the complex numbers. Secondly, if we wish to compute $\text{Gal}(f/\mathbb{Q})$ we may assume that $f(x)$ has rational integer coefficients, for, if not, we may apply the following transformation to $f(x)$:

Let c be the least common multiple of the denominators of the coefficients of $f(x)$. Then

$$g(x) = c^n f(x/c)$$

is a monic polynomial in $\mathbb{Z}[x]$. If (v_1, \dots, v_n) are the zeros of $f(x)$ then (cv_1, \dots, cv_n) are the zeros of $g(x)$, and with respect to these orderings, $\text{Gal}(g/\mathbb{Q}) = \text{Gal}(f/\mathbb{Q})$.

1.2 CONTENT AND CONTRIBUTION OF THIS THESIS

Let $f(x)$ be a separable polynomial in $K[x]$, $n = \deg(f) > 0$.

In this thesis we are concerned with the problem of computing $\text{Gal}(f/K)$ when we have a factorization algorithm for polynomials in $K[x]$. Although there exists a finite algorithm for solving this problem (see Section 2.1), from a

feasible computational viewpoint, finding $\text{Gal}(f/K)$ is difficult.

In this thesis we pay special attention to the case where $K = \mathbb{Q}$ and $f(x)$ is irreducible over K . In this case $\text{Gal}(f/K)$ is transitive (see Proposition 2.7). We note that for reducible $f(x)$ the most efficient methods of finding $\text{Gal}(f/K)$ would probably include determining the intersections of the splitting fields of pairs of irreducible factors of $f(x)$.

We will discuss algorithms which determine invariants of the conjugacy class of $\text{Gal}(f/K)$, when given $f(x)$. The aim is to efficiently determine enough information to specify a representative of the conjugacy class of $\text{Gal}(f/K)$. We use the tables in Appendix 1 of non-conjugate transitive permutation groups (of degree up to 8), and invariants of their respective conjugacy classes. These tables were supplied by G. Butler.

In Chapter 2 we discuss computational methods used to determine invariants of $\text{Gal}(f/K)$, including work done previously. We discuss in detail the use of resolvent polynomials and show how the linear resolvent can be used in determining $\text{Gal}(f/K)$.

In Chapter 3 we describe a new, practical, exact algorithm which uses polynomial resultants to compute linear resolvent polynomials. Our algorithm requires some

restrictions on the base field K when $\text{char}(K) \neq 0$.

In Chapter 4, we implement the algorithm of Chapter 3 over $K = \mathbb{Z}_p$, for p sufficiently large, as a modular algorithm which computes linear resolvents over \mathbb{Z} for monic polynomials $f(x)$ in $\mathbb{Z}[x]$. Also in Chapter 4, we include examples which illustrate methods described in this thesis.

An extension of this work would be to develop practical exact methods to compute an arbitrary resolvent polynomial.

For every transitive permutation group G of degree up to 7 we have computed a polynomial $f(x)$ such that $\text{Gal}(f/\mathbb{Q}) = G$. These polynomials appear in Appendix 2. This is the first such list of which the author is aware. In Appendix 3 we list new examples of degree 7 polynomials with the simple, group $\text{PSL}(3,2)$ as Galois group. These polynomials were found by computer searching as were many other of our examples.

CHAPTER 2

METHODS OF DETERMINING GALOIS GROUPS

In this chapter we discuss algorithms to determine properties of the Galois group of a polynomial. The aim is to determine sufficient properties efficiently to specify the conjugacy class of the Galois group. We include work done previously in this chapter, and our discussion centres on the resolvent polynomial.

For an historical perspective on (computational) Galois theory see [DEH,MAT,FOU-1].

2.1 DETERMINING THE GALOIS GROUP IN FINITELY MANY STEPS

Let $f(x)$ be in $K[x]$, $n = \deg(f) > 0$, furthermore suppose $f(x)$ has distinct zeros, v_1, \dots, v_n , in the splitting field of $f(x)$ over K .

If there is an algorithm for factoring multivariate polynomials over K then one can determine $\text{Gal}(f/K)$ in a finite number of steps using a method detailed in van der Waerden [VDW,p.189]. We note that such a factoring algorithm exists when there is an algorithm for factorizing univariate polynomials over K [VDW,p.135].

This Galois group algorithm proceeds as follows:

Form the expression

$$t = x_1 v_1 + \dots + x_n v_n,$$

where x_1, \dots, x_n are indeterminates. Let $t_1, t_2, \dots, t_n!$ be the distinct expressions obtained from t by applying all the possible permutations to the indices of the x_i . Set

$$F = F(z, x_1, \dots, x_n) = \prod_{i=1}^{n!} (z - t_i).$$

F has coefficients symmetric in the v_i , and hence the coefficients of F can be expressed in terms of the coefficients of $f(x)$ and the x_i . Let the factorization of F into irreducible factors over $K[z, x_1, \dots, x_n]$ be

$$F = F_1 F_2 \dots F_r.$$

The permutations of the x_i which leave invariant any factor, say F_1 , form a group G .

THEOREM 2.1 ([VDW, p.189]) If we assume that the zeros of $f(x)$ are ordered so that $x_1 v_1 + \dots + x_n v_n$ is a zero of F_1 , then $\text{Gal}(f/K) = G$.

This method is clearly impractical from a computational point of view. However, the result of Theorem 2.1 is used to prove [VDW, p.191] a computationally useful result for the case $K = \mathbb{Q}$. This result is stated in Theorem 2.2.

2.2 THE DETERMINATION OF CYCLE TYPES IN $\text{Gal}(f/\mathbb{Q})$

Let $f(x)$ be a monic separable polynomial in $\mathbb{Z}[x]$, $n = \deg(f) > 1$, and let p be a prime.

We define the cycle type of a permutation P in S_n to be the partition of n induced by the lengths of the disjoint cycles of P . The factor type of $f(x) \bmod p$ is defined to be the partition of n induced by the degrees of the irreducible factors of $f(x) \bmod p$ over \mathbb{Z}_p . A useful method to discover information about $\text{Gal}(f/\mathbb{Q})$ is to determine cycle types of permutations in $\text{Gal}(f/\mathbb{Q})$ by factorizing $f(x) \bmod p$ over \mathbb{Z}_p for primes p not dividing $\text{disc}(f)$. This method has been discussed by many authors including van der Waerden [VDW], Zassenhaus [ZAS-2] and McKay [MCK].

THEOREM 2.2. For any prime p not dividing $\text{disc}(f)$, the factor type of $f(x) \bmod p$ is the cycle type of some permutation in $\text{Gal}(f/\mathbb{Q})$.

The following result which follows from the density theorem of Chebotarev may also be used (see [SCH,LAG]).

THEOREM 2.3. Let T be a partition of n . Then as $k \rightarrow \infty$, the proportion of occurrences of T as the factor type of $f(x) \bmod p_i$, $i=1,\dots,k$, (p_1,\dots,p_k distinct primes) tends to the proportion of permutations in $\text{Gal}(f/\mathbb{Q})$ having the cycle type T .

We may factorize $f(x) \bmod p$ over \mathbb{Z}_p using the algorithm of Berlekamp [KNU, p.420-429]. However, as we are only interested in the factor type of $f(x) \bmod p$, we may use the partial factorization method described by Knuth [KNU, p.429-430], which provides us with the necessary information.

Tables 3C, ..., 8C in Appendix 1 contain the distribution of permutation cycle types in transitive permutation groups of degrees 3 to 8 respectively. These tables are used when applying Theorems 2.2 and 2.3. Applying Theorem 2.2, we can determine cycle types of permutations in $\text{Gal}(f/\mathbb{Q})$. After doing this, we exclude permutation groups as candidates for $\text{Gal}(f/\mathbb{Q})$ which do not contain permutations having these determined cycle types. Applying Theorem 2.3, we can make an educated guess as to $\text{Gal}(f/\mathbb{Q})$ after factorizing $f(x) \bmod p$ for a "sufficient" number of primes p . Note, however, that there are two distinct groups of even permutations of degree 8 (T32 and T33) having the same number of elements of each cycle type.

If $\text{Gal}(f/\mathbb{Q}) = S_n$ or A_n then we can usually quickly determine $\text{Gal}(f/\mathbb{Q})$ by applying Theorem 2.2 and using the fact that $\text{Gal}(f/\mathbb{Q}) \leq A_n$ iff $\text{disc}(f)$ is a rational integral square (see Proposition 2.12).

We now give an example of a polynomial having S_6 as Galois group over \mathbb{Q} .

EXAMPLE 2.4. Let $f(x) = x^6 + 2x + 2$; $\text{disc}(f) = -2^6 \cdot 89 \cdot 227$. $f(x)$ is irreducible over \mathbb{Q} using Eisenstein's criterion with the prime 2. The factor type of $f(x) \pmod{7}$ is $(3, 2, 1)$ and the factor type of $f(x) \pmod{11}$ is $(5, 1)$. Hence $\text{Gal}(f/\mathbb{Q})$ is transitive and contains permutations with cycle types $(3, 2, 1)$ and $(5, 1)$. This implies that $\text{Gal}(f/\mathbb{Q}) = S_6$ (see Table 6C in Appendix 1).

We now give an example which shows how useful Theorem 2.3 can be to make an educated guess as to $\text{Gal}(f/\mathbb{Q})$.

EXAMPLE 2.5. Let $f(x) = x^7 - 14x^5 + 56x^3 - 56x + 22$; $\text{disc}(f) = 2^6 \cdot 7^{10}$. $f(x) \pmod{p}$ was factored over \mathbb{Z}_p for the 42 primes p in the interval $[2, 193]$ which do not divide $\text{disc}(f)$. For one prime the factor type is (1^7) , for thirty primes the factor type is $(3^2, 1)$ and for eleven primes it is (7) . Referring to Table 7C in Appendix 1, one feels confident from this information that $\text{Gal}(f/\mathbb{Q}) = 7T3$, the Frobenius group of order 21. In fact one can show that $\text{Gal}(f/\mathbb{Q})$ is indeed $7T3$ (see Section 4.3, Example 4.1). Note that since $\text{disc}(f)$ is a square, $\text{Gal}(f/\mathbb{Q}) \leq A_7$. This, together with the cycle types in $\text{Gal}(f/\mathbb{Q})$ we have determined, has narrowed the candidates for $\text{Gal}(f/\mathbb{Q})$ down to $7T3$, $7T5$, and $7T6 (= A_7)$.

Complex conjugation is an automorphism of any subfield of the complex numbers. If $f(x)$ is separable over \mathbb{Q} , then complex conjugation induces an element in $\text{Gal}(f/\mathbb{Q})$ of cycle type $(2^c, 1^r)$, where c is the number of complex conjugate

pairs of zeros of $f(x)$ and r is the number of real zeros of $f(x)$. The number of real zeros of a polynomial over \mathbb{Q} can be determined by a Sturm polynomial remainder sequence [BUR, vol. 1, p. 198-203]. We note that the polynomial $f(x)$ in Example 2.5 has all zeros real. This is a necessary condition for $|\text{Gal}(f/\mathbb{Q})|$ to be odd.

The preceding factorizations modulo p , discriminants, and the number of real zeros, were calculated using program ONEPOLY written by Regener and Rohlicek.

2.3 THE RESOLVENT POLYNOMIAL

Let $f(x)$ be separable over K , $n = \deg(f) > 0$, and let an ordering of the zeros of $f(x)$ be $\underline{v} = (v_1, \dots, v_n)$. Resolvent polynomials are classical and computationally useful tools to determine $\text{Gal}(f/K)$, and it is the method we concentrate on. For F in $K[x_1, \dots, x_n]$, we use the resolvent polynomial $R(F, f)$ (with distinct zeros) to determine the orbit length partition of F^*S_n under $\text{Gal}(f/K)$.

2.3.1 THEORETICAL DEVELOPMENT

Let N be the splitting field of $f(x)$ over K . Then $G(N/K)$ acts on N in a natural way as a group of automorphisms. We now show that each orbit of elements in N under the action of $G(N/K)$ consists precisely of the zeros of a monic irreducible polynomial over K . First we prove

the following:

LEMMA 2.6. Let $W = \{w_1, \dots, w_k\}$ be contained in N (w_i distinct), and $g(x) = \prod_{i=1}^k (x-w_i)$. Then $G(N/K)$ maps W onto W if and only if $g(x)$ is in $K[x]$.

PROOF. Let $g(x) = \sum_{i=0}^k a_i x^i$, w in W , and S in $G(N/K)$.

Suppose $g(x)$ is in $K[x]$. As S is an automorphism of N fixing K we have:

$$\begin{aligned} 0 &= g(w) = g(w)S = \sum a_i S(w^i)S \\ &= \sum a_i (wS)^i = g(wS). \end{aligned}$$

Thus wS is in W for all w in W and S in $G(N/K)$. Hence $G(N/K)$ maps W onto W .

Conversely, suppose $G(N/K)$ maps W onto W . Then each element S in $G(N/K)$ induces a permutation of W . Thus $a_i S = a_i$ for each coefficient a_i of $g(x)$ because a_i is a symmetric function of w_1, \dots, w_k . This implies that a_i is in K . //

PROPOSITION 2.7. Let $G = G(N/K)$, and w in $W = \{w_1, \dots, w_k\}$ contained in N (w_i distinct). Denote by wG the set $\{wS : S \in G\}$. Then $W = wG$ if and only if $g(x) = \prod_{i=1}^k (x-w_i)$ is an irreducible polynomial over K .

PROOF. If $wG = W$, then by the previous lemma, $g(x)$ is in $K[x]$. Suppose $g(x)$ is reducible. Then $g(x)$ has a factor $h(x)$ in $K[x]$ where $h(x) = \prod_{i \in I} (x-w_i)$, for some I properly contained in $\{1, \dots, k\}$. Then by the previous lemma G maps

$\{w_i : i \text{ in } I\}$ onto itself, which contradicts the fact that $wG = W$.

Conversely suppose that $g(x)$ is a irreducible polynomial in $K[x]$. By the previous lemma, we know that G maps W onto itself. Thus wG is contained in W . Suppose $wG = \{w_i : i \text{ in } I\}$, where I is properly contained in $\{1, \dots, k\}$.

Then by the previous lemma, $h(x) = \prod_{i \in I} (x - w_i)$ is in $K[x]$. Since $h(x)$ is a proper divisor of $g(x)$, we have arrived at the desired contradiction. //

We apply the preceding result to determine the information available from the factorization of a given resolvent polynomial.

Let F be in $K[x_1, \dots, x_n]$. Recall that the resolvent polynomial over K associated with F and $f(x)$ is:

$$R(F, f) = \prod_{i=1}^k (x - F_i(\underline{v}))$$

where $\{F_1, \dots, F_k\} = F^*S_n$ (F_i distinct).

For S in $G(N/K)$, let $S \rightarrow \bar{S}$ under the representation of $G(N/K)$ onto $\text{Gal}(f/K)$ discussed in Section 1.1.2. First we show:

LEMMA 2.8. $F(\underline{v})S = F^*\bar{S}(\underline{v})$.

PROOF. $F(v_1, \dots, v_n)S = F(v_1S, \dots, v_nS)$
 $= F(v_1\bar{S}, \dots, v_n\bar{S}) = F^*\bar{S}(v_1, \dots, v_n)$. //

Thus $\text{Gal}(f/K)$ acts on polynomials in the zeros of $f(x)$ in precisely the same way that $G(N/K)$ does.

PROPOSITION 2.9. Let t be in I contained in $\{1, \dots, k\}$.

- (1) If $F_t * \text{Gal}(f/K) = \{F_i : i \text{ in } I\}$ and the $F_i(\underline{v})$ are distinct for i in I , then

$g(x) = \prod_{i \text{ in } I} (x - F_i(\underline{v}))$ is an irreducible polynomial over K .

- (2) If $g(x) = \prod_{i \text{ in } I} (x - F_i(\underline{v}))$ is a non-repeated irreducible factor of $R(F, f)$ then $F_t * \text{Gal}(f/K) = \{F_i : i \text{ in } I\}$.

PROOF.

- (1) Apply Lemma 2.8 and Proposition 2.7.

- (2) As N is separable over K , $g(x)$ must have distinct zeros.

By Proposition 2.7 and Lemma 2.8, $\{F_i(\underline{v}) : i \text{ in } I\} = \{F^*P(\underline{v}) : P \text{ in } \text{Gal}(f/K)\}$. As $g(x)$ is a non-repeated factor of $R(F, f)$, for all i in I and $j=1, \dots, k$, $F_i(\underline{v}) = F_j(\underline{v})$ if and only if $i=j$. The result follows. //

COROLLARY 2.10. Suppose $R(F, f)$ has distinct zeros.

Then the orbit length partition of F^*S_n under $\text{Gal}(f/K)$ is the same as the partition of $\deg(R(F, f))$ induced by the degrees of the irreducible factors of $R(F, f)$ over K .

A method of dealing with the occurrence of repeated zeros of $R(F, f)$ is the use of an appropriate Tschirnhaus transformation [BUR, vol. 2, p. 171-175] applied to $f(x)$.

Now suppose $R(F, f)$ has distinct zeros:

$F^*P_1(\underline{V}), \dots, F^*P_k(\underline{V})$, where $\{P_1, \dots, P_k\}$ is a set of right coset representatives of $\text{stab}_{S_n}(F)$ in S_n . We see that $\text{Gal}(f/K)$ acts on the zeros of $R(F, f)$ by permuting the P_i . We have $\text{Gal}(R(F, f)/K) = \text{im}(\text{rep}(\text{Gal}(f/K), \underline{P}, *))$, where $\underline{P} = (P_1, \dots, P_k)$ and the action * is defined by $P_i * T = P_i T$ for all T in $\text{Gal}(f/K)$ and i in $\{1, \dots, k\}$. Note that the orbit length partition of F^*S_n under $\text{Gal}(f/K)$ depends only on $\text{stab}_{S_n}(F)$.

The following result is also of interest:

LEMMA 2.11. Let $F_t(\underline{V})$ be a zero of a non-repeated irreducible factor of the resolvent polynomial $R(F, f)$. Then $K(F_t(\underline{V}))$ is the fixed field corresponding to H , where $H \leq G(N/K)$ maps onto $\text{stab}_{\text{Gal}(f/K)}(F_t)$ under $\text{rep}(G(N/K), \underline{V}, *)$.

PROOF. Now $F_t(\underline{V})S \neq F_t(\underline{V})$ for all S in H . If $F_t(\underline{V})S = F_t(\underline{V})$ for some S not in H , then this implies that $F_t(\underline{V})$ is a repeated zero of $R(F, f)$, which is a contradiction. //

2.3.2 CONSTRUCTION AND FACTORIZATION OF RESOLVENTS

The resolvent polynomial $R(F, f)$ can be constructed by expanding $R(F, f)$ symbolically in the zeros of $f(x)$ and then determining the coefficients of $R(F, f)$ as polynomials in the coefficients of $f(x)$. See Lauer [LAU] for methods related to symmetric polynomials. Unfortunately, unless $\deg(R(F, f))$ is small or $f(x)$ is sparse, this leads to very extensive

symbolic manipulation. However, if we use this method, we get an explicit formula for the coefficients of $R(F,f)$ in terms of the coefficients of $f(x)$. Such formulas have been published for specific resolvent polynomials in [BER,DEH,ERB,MAT].

In Chapter 3, we describe a new exact algorithm to construct linear resolvent polynomials. This algorithm does not expand the resolvent symbolically in the zeros of $f(x)$.

For $K = \mathbb{Q}$, monic $f(x)$ in $\mathbb{Z}[x]$, and F in $\mathbb{Z}[x_1, \dots, x_n]$, we note that the coefficients of $R(F,f)$ are algebraic integers and hence rational integers. Thus if we form $R(F,f)$ using numerical approximations to the zeros of $f(x)$ and we know that the accuracy of these approximations is such that the coefficients of $R(F,f)$ are calculated to within an absolute error less than $1/2$, then we can determine the coefficients of $R(F,f)$ exactly by rounding. Stauduhar [STA-1,STA-2] uses this method (see Section 2.3.4).

In Section 4.1 we discuss a modular approach to computing $R(F,f)$ when $f(x)$ and F are as in the preceding paragraph.

We have assumed we have a factorization algorithm over $K[x]$. For $K = \mathbb{Q}$, factorization algorithms are discussed in [KNU,p.431-434,SCH,ZAS-1]. In practice, for $K = \mathbb{Q}$, monic $f(x)$ in $\mathbb{Z}[x]$, and F in $\mathbb{Z}[x_1, \dots, x_n]$, one can determine candidates for factors of $R(F,f)$ by using numerical

approximations to the zeros of $f(x)$.

2.3.3 FUNCTIONS BELONGING TO GROUPS

Let F be in $K[x_1, \dots, x_n]$ and $G = \text{stab}_{S_n}(F)$. We say that F belongs to G . Note that for P in S_n , F^*P belongs to $P^{-1}GP$, and in addition, $F^*P(V)$ is a zero of $R(F, f)$.

Applying Proposition 2.9, we see that if $\text{Gal}(f/K) \leq P^{-1}GP$ for some P in S_n , then $R(F, f)$ has a linear factor. Conversely, if $R(F, f)$ has a non-repeated linear factor then $\text{Gal}(f/K)$ is contained some conjugate of G .

Resolvents where a linear factor determines if $\text{Gal}(f/K)$ is contained in a group of interest are discussed in [BER,FOU-2,LEF,STA-1,STA-2]. Although linear factors are easy to find, the linear factor can give information only about the Galois group's containment in a group and its conjugates. The complete factorization of a well-chosen resolvent polynomial often distinguishes $\text{Gal}(f/K)$ among many possible candidates.

2.3.3.1 THE ALTERNATING FUNCTION

Suppose $\text{char}(K) \neq 2$ and $n > 1$. Then the function

$$D = D(x_1, \dots, x_n) = \prod_{i < j} (x_i - x_j)$$

belongs to A_n , and is called the alternating function. If P in S_n is a odd permutation, then $D^*P = -D$. Thus

$$R(D, f) = x^2 - (D(\underline{v}))^2.$$

If $f(x)$ has distinct zeros, then $R(D, f)$ has distinct zeros and $R(D, f)$ has a linear factor over K if and only if $D(\underline{v})^2 = \text{disc}(f)$ is a square in K . Thus we have proved:

PROPOSITION 2.12. $\text{Gal}(f/K) \leq A_n$ if and only if $\text{disc}(f)$ is a square in K .

2.3.4 THE METHOD OF STAUDUHAR

In [STA-1], and in a condensed version [STA-2], Stauduhar describes an effective method of determining the Galois group over Q of a monic irreducible polynomial $f(x)$ over Z . He describes the implementation of this method for $n = \deg(f)$ up to 8, and supplies tables of information necessary for this implementation. Schnackenberg [SCH] includes a discussion of Stauduhar's method in his thesis which surveys techniques to calculate Galois groups.

Stauduhar proceeds as follows:

Let $\underline{v} = (v_1, \dots, v_n)$ be an ordering of the zeros of $f(x)$ and suppose that with respect to this ordering we know that $\text{Gal}(f/Q) \leq G$. (Initially we know that $\text{Gal}(f/Q) \leq S_n$). If G has no transitive proper subgroups, then $\text{Gal}(f/Q) = G$. Otherwise we check to see if $\text{Gal}(f/Q) \leq H$, for each maximal transitive subgroup H of G .

For H a maximal transitive subgroup of G , we determine if $\text{Gal}(f/Q) \leq P^{-1}HP$ for some P in G . Choose (from a table) a polynomial F in $\mathbb{Z}[x_1, \dots, x_n]$ such that $\text{stab}_G(F) = H$ and consider the factor of $R(F, f)$:

$$R_G(F, f) = \prod_{i=1}^k (x - F_i(\underline{v})),$$

where $F_i = F * P_i$ ($i=1, \dots, k$, F_i distinct), $\{P_i\}$ a set of right coset representatives of H in G (obtained from a table). $\text{Gal}(f/Q) \leq G$ implies that each element in $\text{Gal}(f/Q)$ induces a permutation of the F_i . Hence $R_G(F, f)$ has rational integer coefficients which are determined by expanding $R_G(F, f)$ using high-precision approximations to the zeros of $f(x)$ and then rounding the approximate coefficients of $R_G(F, f)$. If $\text{Gal}(f/Q)$ is contained in some conjugate of H in G , then $R_G(F, f)$ has an integral zero. Conversely, if $R_G(F, f)$ has a non-repeated integral zero, then $\text{Gal}(f/Q)$ is contained in some conjugate of H in G . We test each approximate zero z of $R_G(F, f)$ which appears to be integral to determine if $R_G(F, f)(\text{round}(z)) = 0$. Suppose $R_G(F, f)$ has a non-repeated integral zero, $F * P(\underline{v})$, P in G . Then $\text{Gal}(f/Q) \leq P^{-1}HP$. We may reorder the zeros of $f(x)$ by setting \underline{v} to (v_{1P}, \dots, v_{nP}) , and with respect to this ordering, $\text{Gal}(f/Q) \leq H$.

If $\text{Gal}(f/Q)$ is contained in no maximal transitive subgroup of G , then $\text{Gal}(f/Q) = G$. Otherwise, we have determined that $\text{Gal}(f/Q) \leq H$ with respect to the ordering \underline{v} ,

where H is a maximal transitive subgroup of G . We may then set G to H and repeat the entire process.

In [STA-1] the information available from a quadratic factor of a resolvent polynomial is discussed.

Stauduhar's method is straightforward and practical. However, highly accurate approximations to the zeros of $f(x)$ are necessary, and one must have much tabulated information available. Furthermore a search down the subgroup lattice of S_n is required since if a function F belongs to G , then F is fixed by the elements of any subgroup of G .

2.3.5 THE USE OF LINEAR RESOLVENT POLYNOMIALS

As usual $f(x)$ is a separable polynomial over K , with zeros v_1, \dots, v_n and splitting field N . Let the multiset $M = [e_1, \dots, e_r]$, where $e_i \in K$ and $0 < r \leq n$. We call r the length of M . We may also write

$$M = [a_1^{m_1}, \dots, a_k^{m_k}],$$

where the a_i are distinct and $m_i > 0$ is the multiplicity of a_i in M .

Recall that the linear resolvent polynomial $LR(M, f)$ associated with M and $f(x)$ is the resolvent polynomial $R(F, f)$, where $F = e_1x_1 + \dots + e_rx_r$. We treat any zero elements of M as symbolic placeholders. The degree of $LR(M, f)$ is the number of ways of choosing r objects out of n , times the

number of distinct permutations of the elements of M . Thus

$$(2.1) \deg(LR(M,f)) = \binom{n}{r} r!/(m_1! \dots m_k!) \\ = n!/(m_1! \dots m_k!(n-r)!).$$

Linear resolvents form a general class of useful resolvent polynomials for $f(x)$ of any degree. Often the factorization of linear resolvents of relatively low degree can be used to determine $\text{Gal}(f/K)$, exactly.

2.3.5.1 ACTION ON SETS AND SEQUENCES

A permutation group $G \leq S_n$ acts on the r -sets contained in $\{1, \dots, n\}$ where the action is defined by $\{i_1, \dots, i_r\} * P = \{i_1 P, \dots, i_r P\}$ for all P in G . It is clear that the action of G on $F^* S_n$, where $F = x_1 + \dots + x_r$, is equivalent to the action of G on the r -sets of $\{1, \dots, n\}$. Thus the factorization of $LR([1^r], f)$ (with distinct zeros) determines the orbit length partition of $\{1, \dots, r\} * S_n$ under $\text{Gal}(f/K)$. McKay [MCK], and Erbach, Fischer and McKay [ERB] suggest using resolvents of this form in order to determine the transitivity on r -sets of $\text{Gal}(f/K)$.

The following remark is of interest: Suppose $f(x)$ is irreducible ($\text{Gal}(f/K)$ is transitive) and $n=rs$, s an integer, $s \neq 1, n$. Then $LR([1^r], f)$ (with distinct zeros) has t irreducible factors of degree s if and only if $\text{Gal}(f/K)$ has t systems of imprimitivity of s blocks of size r .

A permutation group $G \leq S_n$ acts on the set of r -sequences (i_1, \dots, i_r) , with i_j in $\{1, \dots, n\}$ and the i_j distinct ($j=1, \dots, r$). This action is defined by $(i_1, \dots, i_r)^*P = (i_{1P}, \dots, i_{rP})$ for all P in G . It is clear that the action of G on F^*S_n , where $F = e_1x_1 + \dots + e_rx_r$, e_i distinct, is equivalent to the action of G on r -sequences.

Now suppose $LR(M, f) = LR([e_1, \dots, e_r], f)$ has distinct zeros and the e_i are distinct. $LR(M, f)$ is reducible iff $\text{Gal}(f/K)$ is not r -ply transitive.

There is also a simple field theoretic interpretation to the factorization of this $LR(M, f)$. Let $z = e_1v_{1P} + \dots + e_rv_{rP}$ be a zero of $LR(M, f)$ (P in S_n). We see that $\text{stab}_{G(N/K)}(z) = \bigcap_{i=1}^r \text{stab}_{G(N/K)}(v_{iP})$, and hence by LEMMA 2.11, $K(z) = K(v_{1P}, \dots, v_{rP})$. The degrees of the irreducible factors of $LR(M, f)$ correspond to the degrees over K of non-conjugate subfields of N generated by r -sets of the zeros of $f(x)$. In particular we note that if $r=2$ and $f(x)$ is irreducible, then $LR(M, f)$ has irreducible factors all of degree n if and only if $N = K(v_i)$ for each zero v_i of $f(x)$, since $K(v_i) = K(v_j)$ for all $i, j = 1, \dots, n$ in this case. We also note that if $r=n$ then $LR(M, f)$ has degree $n!$ and $N = K(z)$ for each zero z of $LR(M, f)$.

Tables 3D to 8D contain the orbit length partitions of r -sets and 2-sequences under the action of the transitive

permutation groups of degrees 3 to 8 respectively. For irreducible $f(x)$, these tables are used to determine candidates for $\text{Gal}(f/K)$ given the factorization of a linear resolvent which determines the orbit lengths of the action of $\text{Gal}(f/K)$ on r-sets or 2-sequences.

2.3.5.2 DIFFERENTIATING ALL TRANSITIVE GROUPS

OF DEGREE UP TO 7

Suppose $\text{char}(K) \neq 2$. If $\text{Gal}(f/K)$ is transitive and we know from $\text{disc}(f)$ whether $\text{Gal}(f/K) \leq A_n$, then for $n=3, 4, 5, 7$, the conjugacy class of $\text{Gal}(f/K)$ is determined completely by the orbit lengths of the action of $\text{Gal}(f/K)$ on 2-sets, 3-sets and 2-sequences, with the exception of distinguishing group 5T3 from 5T5.

Group 5T3 can be distinguished from 5T5 ($= S_5$) in the following way. Let $F = (x_1+x_2-x_3-x_4)^2$ and note that $R(F, f)(x^2) = LR([1^2, -1^2], f)(x)$. We use this linear resolvent to compute $R(F, f)$. For $\deg(f) = 5$, $\deg(R(F, f)) = 15$, and the orbit length partition of $F * S_5$ under 5T3 is (10, 5).

For degree 6, all the transitive groups can be differentiated by $\text{disc}(f)$ and the action on 2-sets, 3-sets and 2-sequences except to distinguish group T8 from T11, T9 from T13, and T14 from T16 (see Table 6D). To distinguish these groups one can use ad hoc techniques, or Stauduhar's

method if $K = \mathbb{Q}$.

We briefly outline a suitable ad hoc technique. We assume that all polynomials discussed have distinct zeros.

Let $D = \text{disc}(f)$ not be a square in K , and $d(x) = x^2 - D$. If we are working over \mathbb{Z} we may take D to be the squarefree part of $\text{disc}(f)$. Let $r(x)$ be a monic irreducible factor over K of a resolvent polynomial $R(F, f)$. Suppose $r(F_t(\underline{V})) = 0$ for some ordering \underline{V} of the zeros of $f(x)$ and F_t in F^*S_n . The following are equivalent:

- (1) $\text{stab}_{\text{Gal}(f/K)}(F_t) \leq A_n$.
- (2) $K(F_t(\underline{V}))$ contains $K(D^{1/2})$.
- (3) $SZ(r(x), d(x))$ has a factor over K of degree $\deg(r)$ (see Section 3.2.5 for an explanation of SZ , and also see [VDW, p. 126-127]).

Now suppose $n = 6$.

Suppose $\text{Gal}(f/K) = T_8$ or T_{11} . Let $r(x)$ be the monic irreducible factor (over K) of degree 12 of $LR([1^3], f)$. Then $\text{Gal}(f/K) = T_8$ if and only if $SZ(r(x), d(x))$ has a factor (over K) of degree 12.

Suppose $\text{Gal}(f/K) = T_9$ or T_{13} . Let $r(x)$ be the monic irreducible factor of degree 2 of $LR([1^3], f)$. Then $\text{Gal}(f/K) = T_9$ if and only if $SZ(r(x), d(x))$ has a factor of degree 2.

Suppose $\text{Gal}(f/K) = T_{14}$ or T_{16} . Let $r(x) = \text{LR}([1^3], f)$.
Then $\text{Gal}(f/K) = T_{14}$ if and only if $\text{SZ}(r(x), d(x))$ has a
factor of degree 20.

CHAPTER 3

LINEAR RESOLVENT POLYNOMIAL CONSTRUCTION

In this chapter we describe an algorithm to construct any linear resolvent polynomial over a field K subject to the restrictions in Section 3.1. The algorithm is exact, uses polynomial resultants and does not expand the resolvent symbolically in the zeros of $f(x)$. This approach was inspired by Trager [TRA], who used polynomial resultants in a similar manner to factorize polynomials over algebraic extension fields.

The usefulness of the linear resolvent in computing $\text{Gal}(f/K)$ when we have a factorization algorithm over $K[x]$ has been discussed in Section 2.3.5.

3.1 RESTRICTIONS ON THE FIELD

The linear resolvent algorithm is designed to work over an arbitrary field K , except for the following restrictions:

If $\text{char}(K) \neq 0$ then we require that $\text{char}(K) > D$, where D is the maximum degree of any polynomial used or constructed by the main algorithm or any sub-algorithm. If $\text{char}(K) \neq 0$, then $\text{char}(K)$ is a prime, and $\text{char}(K) > D$ if and only if $\text{char}(K) \nmid D$.

If K is finite, we need K large enough to construct required polynomials by interpolation. For this requirement, $|K| > 2D$ is sufficient. We note that our interest is not in finding the Galois group of a polynomial over a finite field (such a Galois group is always cyclic), but we may use resolvent polynomials over finite fields in a modular algorithm (see Chapter 4).

3.2 POLYNOMIAL OPERATIONS

In this section we describe our basic operations on polynomials over K . We use these operations for the linear resolvent algorithm.

3.2.1 THE GREATEST COMMON DIVISOR

Let $f = f(x)$, $g = g(x)$ be polynomials in $K[x]$. We assume that $f(x)$ and $g(x)$ are not both the zero polynomial.

DEFINITION 3.1. The greatest common divisor of $f(x)$ and $g(x)$, denoted $\gcd(f, g)$, is defined to be the monic polynomial in $K[x]$ of largest degree dividing both $f(x)$ and $g(x)$.

If $\deg(g) > 0$, by the polynomial division algorithm there exist $q(x)$, $r(x)$ in $K[x]$ such that $f(x) = q(x)g(x) + r(x)$, $0 \leq \deg(r) < \deg(g)$. We denote this $r(x)$ by $f \bmod g$. As any common divisor of f and g divides $f \bmod g$, we may use the following recursive formulation of the gcd to compute

$\text{gcd}(f, g)$:

If $g(x)$ is the zero polynomial,

then $\text{gcd}(f, g) = f(x)/(\text{leading coefficient of } f(x))$;

else, if $\deg(g) = 0$, then $\text{gcd}(f, g) = 1$;

else, $\text{gcd}(f, g) = \text{gcd}(g, f \bmod g)$.

Let e be a non-negative integer, and let N be the splitting field of $f(x)$ over K . We say that $f(x)$ has a zero v of multiplicity e , if $(x-v)^e \mid |f(x)$ in $N[x]$. We write $e = \text{mult}(v, f)$.

We note that $\text{gcd}(f, g)$ over any extension L of K is the same as $\text{gcd}(f, g)$ over K . This is because the gcd calculation is carried out exactly the same way over L . In particular, for L the splitting field of $f(x)g(x)$, the zeros of $\text{gcd}(f, g)$ are the common zeros of f and g , and if v is a zero of $\text{gcd}(f, g)$, then $\text{mult}(v, \text{gcd}(f, g)) = \min\{\text{mult}(v, f), \text{mult}(v, g)\}$.

3.2.2 THE RESULTANT

Let $f = f(x)$, $g = g(x)$ be polynomials in $K[x]$. Let $f(x) = a(x-v_1)\dots(x-v_n)$ and $g(x) = b(x-w_1)\dots(x-w_m)$ over the splitting field of $f(x)g(x)$. Furthermore assume that $n = \deg(f) > 0$, and $m = \deg(g)$.

We treat the resultant in a similar manner as Childs [CHI, p.283]. See also Collins [COL].

DEFINITION 3.2. The resultant of $f(x)$ and $g(x)$,

$$\text{res}(f,g) = a^m b^n \prod_{i=1}^n \prod_{j=1}^m (v_i - w_j).$$

The resultant is a symmetric function of both the v_i and w_j , and hence $\text{res}(f,g)$ is an element of K . The following facts are immediate consequences of Definition 3.2.

$$(1) \text{res}(f,g) = (-1)^{mn} \text{res}(g,f).$$

$$(2) \text{res}(f,g) = a^m \prod_{i=1}^n g(v_i).$$

(3) If $m = 0$, then $\text{res}(f,g) = b^n$. (For our purposes it is convenient to assume the degree of the zero polynomial is zero, so that here we do not exclude the possibility that $b = 0$.)

We use (1) and (2) to prove the following lemma.

LEMMA 3.3. Suppose $m > 0$, and let $r(x) = f \bmod g$. Then $\text{res}(f,g) = (-1)^{mn} b^{n-\deg(r)} \text{res}(g,r)$.

$$\text{PROOF. } \text{res}(f,g) = (-1)^{mn} \text{res}(g,f)$$

$$= (-1)^{mn} b^n \prod_{i=1}^m (g(w_i)q(w_i) + r(w_i))$$

$$= (-1)^{mn} b^n \prod_{i=1}^m r(w_i)$$

$$= (-1)^{mn} b^{n-\deg(r)} \text{res}(g,r). //$$

Combining (3) and Lemma 3.3, we have a recursive formulation of $\text{res}(f,g)$ similar to the recursive formulation of $\text{gcd}(f,g)$. This formulation is used to compute $\text{res}(f,g)$ efficiently. One can also compute the resultant or gcd

non-recursively by using a polynomial remainder sequence.

3.2.3 THE FORMAL DERIVATIVE AND ITS ZEROS

The formal derivative of a polynomial over a field K is similar to the usual derivative of a real polynomial, and shares many common properties.

DEFINITION 3.4. Let $f(x) = \sum_{i=0}^n a_i x^i$ be a polynomial over K . We define the formal derivative of $f(x)$, denoted f' or $f'(x)$, by

$$f' = f'(x) = \sum_{i=1}^n i a_i x^{i-1},$$

where $i a_i$ means $a_i + \dots + a_i$ (i times).

There is an important relationship between the multiplicity of zeros of $f(x)$ and the zeros of $f'(x)$, which we state in the following proposition.

PROPOSITION 3.5. Suppose $f(x)$ has a zero v of multiplicity $e > 0$. Then if $\text{char}(K) \neq e$, $\text{mult}(v, f') = e-1$.

PROOF. Let $f(x) = (x-v)^e h(x)$.

Then $f'(x) = e(x-v)^{e-1} h(x) + (x-v)^e h'(x)$. Thus $\text{mult}(v, f') \geq e-1$. Now if $(x-v)^e | f'(x)$, then $(x-v) | e h(x)$. This cannot happen as $\text{char}(K) \neq e$ implies that $e \neq 0$ and by the definition of multiplicity, $x-v$ cannot divide $h(x)$. //

COROLLARY 3.6. Suppose $\text{char}(K) \neq n$. For each zero v of $f(x)$ of multiplicity $e > 1$, v is a zero of $\text{gcd}(f, f')$ of

multiplicity $e=1$, and $\gcd(f, f')$ has no other zeros.

3.2.4 "MULTIPLY ZEROS"

Let $f(x)$ be a monic polynomial over K , $n = \deg(f)$, and let the zeros of $f(x)$ be v_1, \dots, v_n . Let d be an element of K . We want to calculate a monic polynomial of degree n having the zeros dv_1, \dots, dv_n . The required polynomial is denoted $MZ(d, f)$ (Multiply Zeros) and is computed as follows:

$$MZ(d, f) = d^n f(x/d), \text{ if } d \neq 0; x^n, \text{ if } d=0.$$

3.2.5 "SUM ZEROS"

Let $f = f(x)$, $g = g(x)$ be monic polynomials in $K[x]$. Let $f(x) = (x-v_1) \dots (x-v_n)$ and $g(x) = (x-w_1) \dots (x-w_m)$ over the splitting field of $f(x)g(x)$.

We need to calculate the monic polynomial in $K[x]$ of degree mn with zeros $v_i + w_j$, ($i=1, \dots, n$, $j=1, \dots, m$). This polynomial is denoted by $SZ(f, g)$ (Sum Zeros) and we note that equality (3.1) holds as the left-hand side and the right-hand side are both degree mn monic polynomials having the same zeros.

$$(3.1) \quad SZ(f, g) = \prod_{i=1}^n g(x-v_i).$$

Thus for any element y in K we know the value of $SZ(f, g)(y)$.

It is:

$$(3.2) \quad \text{SZ}(f,g)(y) = \text{res}(f(x), g(y-x)).$$

If K is sufficiently large (and we assume it is), we can calculate $z_i = \text{SZ}(f,g)(y_i)$, using (3.2), for $i=1,2,\dots,mn+1$ and y_i in K distinct. Then we can determine $\text{SZ}(f,g)$ by interpolation. That is, we find the polynomial $t(x)$ ($= \text{SZ}(f,g)$) of degree at most mn such that $t(y_i) = z_i$, for $i=1,2,\dots,mn+1$. For interpolation algorithms, see [KNU, COL].

3.2.6 "POLYNOMIAL ROOT"

Finally, we need an algorithm to solve the following problem. Let k be a positive integer and let $u(x)$ be a monic polynomial in $K[x]$, $\deg(u) > 0$. Suppose we know that $u(x) = r(x)^k$ for some unknown monic $r(x)$ in $K[x]$. Denote this unique $r(x)$ by $\text{PR}(k,u)$ (Polynomial Root). We compute $\text{PR}(k,u)$ using the algorithm POLYROOT, which follows. We assume $\text{char}(K) > \deg(u)$ or $\text{char}(K) = 0$.

Algorithm POLYROOT;

Input: positive integer k and monic polynomial $u(x)$ in $K[x]$,
 $\deg(u) > 0$, such that $u(x) = r(x)^k$ for some unknown monic
 $r(x)$ in $K[x]$. We assume $\text{char}(K) > \deg(u)$ or $\text{char}(K) = 0$.

Returns : $\text{PR}(k, u)$. ($= r(x)$).

(1) if $k=1$ then $\text{return}(u(x))$, and stop.

(2) set $t(x) \leftarrow u(x)/\text{gcd}(u, u')$;

{ u' is the formal derivative of $u(x)$. $t(x)$ is separable, and
the zeros of $t(x)$ are precisely the distinct zeros of
 $u(x)$ (recall Corollary 3.6).}

(3) set $r(x) \leftarrow t(x)$, and $s(x) \leftarrow u(x)$;

(4) while $\deg(r) < \deg(u)/k$, execute steps (4.1), ..., (4.3);

(4.1) set $s(x) \leftarrow s(x)/t(x)^k$;

(4.2) set $t(x) \leftarrow \text{gcd}(s, t)$;

{In the i -th iteration of this loop, at this point, the
zeros of $t(x)$ are precisely the distinct zeros v of
 $u(x)$ such that $\text{mult}(v, u) > i$.}

(4.3) set $r(x) \leftarrow t(x)r(x)$;

(5) $\text{return}(r(x))$, and stop.

3.3 MULTISSET OPERATIONS

We define the operations + and - for multisets. They
are similar respectively to union and difference for sets,
except that multiplicities are counted. We use these
operations in the proof following and in the linear
resolvent algorithm. $\text{mult}(e, M)$ denotes the multiplicity of

the element e in the multiset M .

Let M and N be multisets and let e an element of the "universal" set from which M and N draw their elements.

Then $M + N$ is a multiset, and $\text{mult}(e, M + N) = \text{mult}(e, M) + \text{mult}(e, N)$. $M - N$ is a multiset and $\text{mult}(e, M - N) = \text{mult}(e, M) - \text{mult}(e, N)$ if $\text{mult}(e, M) > \text{mult}(e, N)$; and $\text{mult}(e, M - N) = 0$ otherwise.

3.4 CONSTRUCTIVE PROOF

Let K be a field satisfying the restrictions described in Section 3.1. Let $f(x)$ be a monic polynomial in $K[x]$, $n = \deg(f) > 0$, and let the zeros of $f(x)$ be v_1, \dots, v_n . Let e_1, \dots, e_r be in K , $0 < r \leq n$, and let $M = [e_1, \dots, e_r]$. We now prove:

PROPOSITION 3.7. The linear resolvent polynomial $\text{LR}(M, f)$ can be constructed over K using only the operations MZ , SZ , and PR discussed in Section 3.2.

PROOF. By induction on r , the length of M .

If $r = 1$ then $\text{LR}(M, f) = \text{MZ}(e_1, f)$.

Now suppose $r > 1$. Let $\bar{M} = [e_1, \dots, e_{r-1}] = [a_1^{m_1}, \dots, a_k^{m_k}]$, where a_1, \dots, a_k are distinct and $m_i = \text{mult}(a_i, \bar{M}) > 0$ for $i=1, \dots, k$. By the inductive hypothesis we can compute

$$t(x) = \text{SZ}(\text{LR}(\bar{M}, f), \text{MZ}(e_r, f)).$$

For each zero w of $\text{LR}(\bar{M}, f)$, $t(x)$ has precisely the zeros $w+e_r v_1, \dots, w+e_r v_n$. Thus we see that

$$t(x) = \left(\prod_{i=1}^k \text{LR}(M_i, f)^{c_i} \right) \text{LR}(M, f)^c,$$

where $M_i = (\bar{M} - [a_i]) + [a_i + e_r]$,

$c_i = m_i \deg(\text{LR}(\bar{M}, f)) / \deg(\text{LR}(M_i, f))$,

and $c = (n-r+1) \deg(\text{LR}(\bar{M}, f)) / \deg(\text{LR}(M, f))$.

We can compute c_i and c using the expression (2.1) for the degree of a linear resolvent polynomial. In fact, by straightforward calculation involving these expressions, one sees that $c_i = \text{mult}(a_i + e_r, M_i)$ and that $c = \text{mult}(e_r, M)$.

By hypothesis we can construct

$$s(x) = \prod_{i=1}^k \text{LR}(M_i, f)^{c_i}.$$

Then the desired linear resolvent polynomial can be computed by:

$$\text{LR}(M, f) = \text{PR}(c, t(x)/s(x)), //$$

3.5 ALGORITHM LINRESOLV

Let K be a field satisfying the restrictions stated in Section 3.1. Let $f(x)$ be a monic polynomial in $K[x]$, $n = \deg(f) > 0$. Let e_1, \dots, e_r be in K , $0 < r \leq n$, and let the multiset $M = [e_1, \dots, e_r]$.

The inductive proof of the preceding section motivates our recursive algorithm to construct $LR(M, f)$, the linear resolvent associated with M and $f(x)$. Changes from the method of the proof have been made for considerations of efficiency. The algorithm is called LINRESOLV and is detailed below.

Algorithm LINRESOLV;

Input: a monic polynomial $f(x)$ in $K[x]$ of degree $n > 0$, and a multiset $M = [e_1, \dots, e_r]$, $0 \leq r \leq n$, where $e_1, \dots, e_r \in K$.

We assume K satisfies the restrictions stated in Section 3.1.

Output: $LR(M, f)$, the linear resolvent polynomial associated with M and $f(x)$.

(1) {If any of the elements in M equals 0 (the additive identity of K) then these zeros are significant as symbolic place holders. However, this step allows $LR(M, f)$ to be calculated by considering just the maximal submultiset of M which contains only non-zero elements.}

(1.1) set $m \leftarrow \text{mult}(0, M)$;

(1.2) if $m = 0$ then go to step (2);

(1.3) if $m = r$ then set $t(x) \leftarrow "x"$,

and go to step (1.6);

(1.4) set $\bar{M} \leftarrow M - [0^m]$;

(1.5) set $t(x) \leftarrow LR(\bar{M}, f)$;

{recursive application of this algorithm}

- (1.6) set $d \leftarrow \frac{n-r+m}{m}$, return($t(x)^d$), and stop.
- (2) if $r = 1$ then return($MZ(e_1, f)$), and stop.
- (3) Arrange the elements of M so that $\text{mult}(e_r, M) \leq \text{mult}(e_i, M)$,
for $i=1, \dots, r$;
{This ensures that the degree of the polynomial constructed
in step (4.2) is as small as possible.}

(4)

- (4.1) set $\bar{M} \leftarrow [e_1, \dots, e_{r-1}]$
 $= [a_1^{m_1}, \dots, a_k^{m_k}]$, where a_1, \dots, a_k are distinct and $m_i > 0$ for $i=1, \dots, k$);
- (4.2) set $u(x) \leftarrow LR(\bar{M}, f)$;

{using this algorithm recursively}

- (5) set $s(x) \leftarrow \prod_{i=1}^k LR(M_i, f)^{c_i}$
(where $M_i = (\bar{M} - [a_i]) + [a_i + e_r]$, and $c_i = \text{mult}(a_i + e_r, M_i)$);

{using this algorithm recursively}

(6)

- (6.1) set $c \leftarrow \text{mult}(e_r, M)$, $g(x) \leftarrow MZ(e_r, f)$;
- (6.2) set d to be a positive integer such that for all $b = a^c$, for some a in K ,
- (i) a^d is unique in K , for all solutions a in K of
 $b = a^c$, and
- (ii) we can efficiently compute this a^d .

{We may always take $d=c$. However, it is most efficient to
choose d as small as possible. For example, when $K = Q$:
if c is odd then let $d=1$, and a^d is the unique c -th root}

in \mathbb{Q} of b ; if c is even then let $d=2$, and a^d is the unique positive $(c/2)$ -th root in \mathbb{Q} of b .}

(6.3) set $m \leftarrow d(\deg(u)\deg(g)-\deg(s))/c+1$;
 $\{m = \deg(LR(M,f))^d)+1\}$

(6.4) for m distinct y_i in K ($i=1, \dots, m$) such that
 $s(y_i) \neq 0$, set $z_i \leftarrow \text{res}(u(x), g(y_i-x))/s(y_i)$;
{This is where we need to assume that $|K|$ is "large enough"}

$$\{z_i = SZ(u,g)(y_i)/s(y_i) = (LR(M,f)(y_i))^c\}$$

(6.5) For each z_i , we know that $z_i = a_i^c$ for some
 a_i in K .

$$\{a_i = LR(M,f)(y_i)\}$$

For $i=1, \dots, m$ set $z_i \leftarrow a_i^d$;

{We can do this due to the choice of d as explained in step

(6.2).}

(7) set $t(x)$ to be the polynomial of degree $m-1$ such that

$$t(y_i) = z_i, \text{ for } i=1, \dots, m;$$

{using an interpolation algorithm}

(8) return(PR(d, t)), and stop.

3.6 REMARKS

As r increases, the efficiency of Algorithm LINRESOLV decreases markedly. However in practice, r is usually quite small; often $r \leq 3$. For a given field K , empirical observations can be made to determine the practical range for r and n . For example, using the implementation

described in Chapter 4 over $K = \mathbb{Z}_{10000139}$ and $f(x)$ a degree 11 polynomial with no zero coefficients, to compute $\text{LR}([1^r], f)$ for $r=2, 3, 4, 5$, it took respectively 3, 13, 129, 426 CPU seconds.

When Algorithm LINRESOLV computes one resolvent polynomial, it must usually compute other resolvents recursively. If these "byproduct" resolvents are useful they should be saved. For example, to compute $\text{LR}([1^3], f)$, LINRESOLV must also compute $\text{LR}([1^2], f)$ and $\text{LR}([1, 2], f)$.

CHAPTER 4

IMPLEMENTATION AND EXAMPLES

Throughout this chapter the following holds:

$f(x) = x^n + \sum_{i=1}^n a_i x^{n-i}$ is in $\mathbb{Z}[x]$, with zeros v_1, \dots, v_n . $M = [e_1, \dots, e_r]$, with e_i in \mathbb{Z} and $0 < r \leq n$,

We discuss our modular algorithm to compute $LR(M, f)$, and the computer implementation of this algorithm. We give examples of the determination of Galois groups over \mathbb{Q} , using this implementation.

4.1 A MODULAR APPROACH TO COMPUTING RESOLVENTS

Let $S(x_1, \dots, x_n)$ be a symmetric polynomial over \mathbb{Z} . By the Fundamental Theorem on Symmetric Polynomials,

$S = T(s_1, \dots, s_n)$, for a unique T in $\mathbb{Z}[x_1, \dots, x_n]$ and s_i is the i -th elementary symmetric polynomial.

Let $f(x) \bmod p = x^n + \sum_{i=1}^n \bar{a}_i x^{n-i}$ have zeros $\bar{v}_1, \dots, \bar{v}_n$, and \bar{S}, \bar{T} be respectively $S \bmod p$, $T \bmod p$. Then over \mathbb{Z}_p :

$$\bar{S}(\bar{v}_1, \dots, \bar{v}_n) = \bar{T}(-\bar{a}_1, \bar{a}_2, \dots, (-1)^n \bar{a}_n).$$

Thus

$$(4.1) \quad S(v_1, \dots, v_n) \bmod p = \bar{S}(\bar{v}_1, \dots, \bar{v}_n).$$

We see that for any F in $\mathbb{Z}[x_1, \dots, x_n]$ such that
 $\text{stab}_{S_n}(F) = \text{stab}_{S_n}(F \bmod p)$:

$$(4.2) \quad R(F, f) \bmod p = R(F \bmod p, f \bmod p),$$

where the latter resolvent is calculated over \mathbb{Z}_p . To compute $R(F, f)$ over \mathbb{Z} , we can compute $R(F, f) \bmod p_i$

(using (4.2)) for distinct primes p_i such that $\prod p_i > 2C$, where C is an upper bound on the magnitude of the coefficients of $R(F, f)$. $R(F, f)$ is then built up over \mathbb{Z} using the Chinese Remainder Algorithm [KNU, p.268-276].

We calculate C by bounding the magnitude of the zeros of $f(x)$, which allows us to calculate a bound on the magnitude of the zeros of $R(F, f)$. If B is an upper bound on the magnitude of the zeros of $R(F, f)$ and $d = \deg(R(F, f))$, then

$$C = \max\{(\frac{d}{i})B^i : 1 \leq i \leq d\}$$

is an upper bound on the magnitude of the coefficients of $R(F, f)$.

4.1.1 BOUNDING THE ZEROS OF $f(x)$

We need to compute a bound A such that $A \geq |v_i|$ for all zeros v_i of $f(x)$. Zassenhaus [ZAS-1] suggested

$$A = \max\{|a_i/(i)|^{1/i}/(2^{1/n}-1) : 1 \leq i \leq n\}.$$

We suggest the following method of computing a suitable bound. Let $g(x) = x^n - \sum_{i=1}^n |a_i| x^{n-i}$, and let $R > 0$ be a strict upper bound on the magnitude of the real zeros of $g(x)$. (By Descartes rule of signs, $g(x)$ has at most one positive real zero (counting multiplicities), so we may take R to be the least positive integer such that $g(R) > 0$.) Now note that for any complex number z such that $|z| \geq R$:

$$|f(z)| \geq g(|z|) \geq g(R) > 0.$$

Thus $R > |v_i|$ for each zero v_i of $f(x)$.

We have found this bound to be often much better than that of Zassenhaus, and we use our bound for the examples of Section 4.3.

4.2 THE IMPLEMENTATION

We have programmed algorithm LINRESOLV over $K = \mathbb{Z}_p$, in the language PASCAL on the Concordia University CDC Cyber 170-800 computer. This program is used to compute $\text{LR}(M, f) \bmod p_i$ for distinct primes p_i . $\text{LR}(M, f)$ is built up over \mathbb{Z} using the Chinese Remainder Algorithm, and then $\text{LR}(M, f)$ is factorized using Hensel factorization [KNU, SCH, ZAS-1]. For these two operations we use programs written by D. Ford in the language ALGEB, on the PDP-11/34 computer. The language ALGEB was designed by D. Ford, and it allows computation with

integers of arbitrary size.

4.2.1 LINRESOLV OVER $K = \mathbb{Z}_p$

\mathbb{Z}_p satisfies the restrictions of Section 3.1 if $p > 2D$, where D is the maximum degree of any polynomial used in the program. In practice we choose p such that p^2 is nearly equal to the largest integer we can operate on ($10^7 < p < 2^{24}$ in our implementation).

Addition, subtraction, and multiplication over \mathbb{Z}_p is implemented by doing these operations over the integers and then applying the PASCAL mod operator to the result. To divide we need multiplicative inverses in \mathbb{Z}_p . Given a in \mathbb{Z} , p/a , we need to determine b in \mathbb{Z} such that $ab \bmod p = 1$. We know

$$1 = \gcd(a, p) = ab + tp,$$

for some b, t in \mathbb{Z} . b can be computed using the (half) extended Euclidean algorithm [KNU, p.325].

The main problem which is dependent on the base field in the implementation of LINRESOLV is the choice of d in step (6.2). Using the notation of step (6) of Algorithm LINRESOLV, we claim that $d = \gcd(c, p-1)$ is appropriate. Note that

$$d = \gcd(c, p-1) = sc + t(p-1),$$

for some s, t in \mathbb{Z} . Let $b = a^c$ for some a in \mathbb{Z}_p . Then

$$b^s = a^{cs} = a^d.$$

This is because $y^{p-1} = 1$, for any $y \neq 0$ in \mathbb{Z}_p . Now we need to show that $y^d = z^d$ for any y, z in \mathbb{Z}_p such that $b = y^c = z^c$. We have

$$y^c = z^c \Rightarrow y^{cs} = z^{cs} \Rightarrow y^d = z^d.$$

When we use algorithm LINRESOLV over \mathbb{Z}_{p_i} in order to compute $LR(M, f)$ over \mathbb{Z} we choose primes p_i such that $j \nmid (p_i - 1)$ for $j = 3, \dots, n$. In this case d is never greater than 2 in step (6.2).

4.3 EXAMPLES

EXAMPLE 4.1. Let $f(x) = x^7 - 14x^5 + 56x^3 - 56x + 22$ (discussed in Example 2.5). We compute and factorize $L(x) = LR([1^3], f)$ of degree 35 to prove that $\text{Gal}(f/\mathbb{Q})$ is group $7T3$, the Frobenius group of order 21.

An upper bound on the magnitude of the zeros of $f(x)$ is 5, and hence 15 is an upper bound on the magnitude of the zeros of $L(x)$. An upper bound on the magnitude of the coefficients of $L(x)$ is $(1/2)10^{42}$. $L(x) \bmod p_i$ is computed for six primes $p_i > 10^7$. This step requires 10 CPU seconds on the CDC Cyber. $L(x)$ is constructed over \mathbb{Z} using the Chinese Remainder Algorithm. Factorizing

$L(x)$ into irreducible factors over \mathbb{Q} , we find $L(x) =$

$L_1(x)L_2(x)L_3(x)$, where

$$L_1(x) = x^7 - 28x^5 + 224x^3 - 448x + 94,$$

$$L_2(x) = x^7 - 28x^5 + 224x^3 - 448x + 192, \text{ and}$$

$$\begin{aligned} L_3(x) &= x^{21} - 84x^{19} + 2436x^{17} - 31136x^{15} + 6358x^{14} \\ &+ 203840x^{13} - 84392x^{12} - 733824x^{11} + 420728x^{10} + 1480192x^9 \\ &- 988064x^8 - 1652036x^7 + 1138368x^6 + 986496x^5 - 620928x^4 \\ &- 284032x^3 + 137984x^2 + 27104x - 10648. \end{aligned}$$

This factorization takes 12 minutes of CPU time on the PDP-11/34.

$L(x)$ has distinct zeros and its factorization shows that the orbit length partition of 3-sets under $\text{Gal}(f/\mathbb{Q})$ is $(7^2, 21)$. From Table 7D in Appendix 1, we see that $\text{Gal}(f/\mathbb{Q})$ is $7T3$.

EXAMPLE 4.2. Let $f(x) = x^5 + 15x + 12$; $\text{disc}(f) = 2^{10}3^45^5$. $f(x)$ is irreducible over \mathbb{Q} , and since $\text{disc}(f)$ is not a square, from Table 5A in Appendix 1 we see that $\text{Gal}(f/\mathbb{Q}) = 5T3$ (the Frobenius group of order 20) or $5T5(S_5)$.

Let $F = (x_1 + x_2 - x_3 - x_4)^2$. We compute and factorize $R(x) = R(F, f)$, of degree 15 to distinguish between the two candidates for $\text{Gal}(f/\mathbb{Q})$ ($R(F, f)(x^2) = LR([1^2, -1^2], f)(x)$; see Section 2.3.5.2).

An upper bound on the magnitude of the zeros of $f(x)$ is 3, and hence 144 is an upper bound on the magnitude of the zeros of $R(x)$. An upper bound on the magnitude of the coefficients of $R(x)$ is 10^{33} .

$R(x) \bmod p_i$ is computed for five primes $p_i > 10^7$. This step requires 11 CPU seconds on the CDC Cyber. $R(x)$ is constructed over \mathbb{Z} using the Chinese Remainder Algorithm. Factoring $R(x)$ into irreducible factors, we find $R(x) = R_1(x)R_2(x)$, where $\deg(R_1) = 5$ and $\deg(R_2) = 10$.

This factorization takes 2 minutes of CPU time on the PDP-11/34.

$R(x)$ has distinct zeros and its factorization shows that $\text{Gal}(f/\mathbb{Q})$ acts intransitively on F^*S_5 , and hence $\text{Gal}(f/\mathbb{Q})$ is $5T3$.

REFERENCES

- [BER] Berwick, W.E.H., On soluble sextic equations,
Proc. London Math. Soc. (2) 29 (1929), 1-28.
- [BIR] Birkhoff, G. and MacLane, S., "A Survey of Modern
Algebra", Macmillan, New York, 1965.
- [BUR] Burnside, W.S. and Panton, A.W., "The Theory of
Equations", vol. 1 and vol. 2, reprint, Dover, New
York, 1960.
- [CHI] Childs, L., "A Concrete Introduction to Higher
Algebra", Springer-Verlag, New York, 1979.
- [COL] Collins, G.E., The calculation of multivariate
polynomial resultants, JACM, 18 (1971), 515-532..
- [DEH] Dehn, E., "Algebraic Equations", reprint, Dover, New
York, 1960.
- [ERB] Erbach, D.W., Fischer, J. and McKay, J., Polynomials
with $PSL(2,7)$ as Galois group, J. Number Theory, 11
(1979), 69-75.
- [FOU-1] Foulkes, H.O., The algebraic solution of equations,
Science Progress, 26 (1931-2), 601-608.
- [FOU-2] Foulkes, H.O., The resolvents of an equation of the
seventh degree, Quart. J. of Math., 2 (1931), 9-19.
- [KNU] Knuth, D.E., "The Art of Computer Programming", vol
2., 2-nd ed., Addison-Wesley, Reading, Mass., 1981.
- [LAG] Lagarias, J.C. and Odlyzko, A.M., Effective versions
of the Chebotarev density theorem, "Algebraic Number
Fields (L-functions and Galois Theory)", Frolich,
A., ed., Academic Press, 1977, 409-464.
- [LAM] LaMacchia, S.E., Polynomials with Galois group
 $PSL(2,7)$, Communications in Algebra, 8 (1980),
983-992.
- [LAU] Lauer, E., Algorithms for symmetrical polynomials,
Proc. 1976 ACM Symp. on Symbolic and Algebraic
Comput., New York, 242-247.
- [LEF] Lefton, P., Galois resolvents of permutation groups,
Amer. Math. Monthly, 84 (1977), 642-644.
- [LON] Long, C.T., "Elementary Introduction to Number

- Theory", Heath, Lexington, Mass., 1972.
- [MCK] McKay, J., Some remarks on computing Galois groups, SIAM J. Comput., 8 (1979), 344-347.
- [MAT] Mathews, G.B., "Algebraic Equations", Cambridge University Press, London, 1930.
- [NEU] Neumann, P.M., Stoy, G.A. and Thompson, E.C., "Groups and Geometry", vol. 1, The Mathematical Institute, Oxford, 1980.
- [SCH] Schnackenberg, D., Calculation of Galois groups, Master's thesis, University of North Dakota, 1978.
- [SHA] Shafarevich, I.R., Construction of fields of algebraic numbers with given solvable Galois group, Izv. Akad. Nauk. SSSR. Ser. Mat., 18 (1954), 525-578.
- [STA-1] Stauduhar, R.P., The automatic determination of Galois groups, Ph.D. Dissertation, University of California, Berkeley.
- [STA-2] Stauduhar, R.P., The determination of Galois groups, Math. Comput., 27 (1973), 981-996.
- [TRA] Trager, B.M., Algebraic factoring and rational function integration, Proc. 1976 ACM Symp. on Symbolic and Algebraic Comput., New York, 219-226.
- [TRI] Trinks, W., Ein Beispiel eines Zahlkörpers mit der Galoisgruppe $PSL(3,2)$ über \mathbb{Q} , Manuscript, Universität Karlsruhe, 1968.
- [VDW] van der Waerden, B.L., "Modern Algebra", vol. 1, tr. Blum, F., Ungar, New York, 1953.
- [ZAS-1] Zassenhaus, H., On Hensel factorization. I, J. Number Theory, 1 (1969), 291-311.
- [ZAS-2] Zassenhaus, H., On the group of an equation, "Computers in Algebra and Number Theory", Birkhoff, G. and Hall, M., eds., SIAM and AMS Proc., 1971, 69-88.

APPENDIX 1

TABLES OF TRANSITIVE GROUPS OF DEGREE UP TO 8
(SUPPLIED BY G. BUTLER)

For each degree we present the information about the transitive groups of that degree in a set of tables. The groups are named, T_1 , T_2 , etc..., for convenience, and if there may be confusion about the degree of the group we write nT_i to mean the i -th group of degree n .

In Table A we list the order of the group, whether it contains only even permutations, the number of inequivalent minimal sets of imprimitivity of each possible type, and the number of conjugacy classes of elements. If the group has a faithful representation of smaller degree this is given in the column headed 'Other Representation', and if the group is known by a common name this name is given in the column headed 'Name'.

In Table B we give a set of generators for each group.

Table C sets out the number of elements of each group with each cycle type.

Table D gives the orbit length partitions of r -sets and 2-sequences (with distinct elements) under the action of each group.

The notation for the group names is as follows: n denotes the cyclic group of order n ; p^n denotes an elementary abelian group of order p^n , where p is a prime; D_n denotes the dihedral group of order n ; Q_8 is the quaternion group of order 8; A_n is the alternating group of degree n ;

Σ_n is the symmetric group of degree n. If A and B are names for groups then $A \cdot B$ denotes a group with a normal subgroup isomorphic to A such that $(A \cdot B)/A$ is isomorphic to B; while $A \times B$ denotes the direct product.

Table 3A: groups of degree 3

Group	Order	Even	Number of classes	Name
T1	3	+	3	A_3
T2	6		3	Σ_3

Table 3B: group generators

$$a = (1, 2, 3)$$

$$T1 = \langle a \rangle$$

$$b = (1, 2)$$

$$T2 = \langle a, b \rangle$$

Table 3C: cycle type distribution

	1^3	2'	3
T1	1	2	
T2	1	3	2

Table 3D

Orbit length partitions of sets and sequences under G

G	2-sets	2-sequences
$\underline{G \triangle A_3}$		
T1	3	3^2

G	2-sets	2-sequences
$\underline{G \triangle A_3}$		
T2	3	6

Table 4A: groups of degree 4

Group	Order	Even	Imprimitive [2 ²]	Number of classes	Name
T1	4		✓	4	4
T2	4	+	✓	4	2 ²
T3	8		✓	5	D ₈
T4	12	+		4	A ₄
T5	24			5	S ₄

Table 4B: group generators

$$a = (1, 3, 4) \quad c = (2, 4)$$

$$b = (1, 3) \quad d = (1, 2)(3, 4)$$

$$T1 = \langle ac \rangle \quad T4 = \langle a, d \rangle$$

$$T2 = \langle bc, d \rangle \quad T5 = \langle ac, d \rangle$$

$$T3 = \langle ac, bc \rangle$$

Table 4C: cycle type distribution

	1 ⁴	1 ²	2 ²	3	4
T1	1	.	1	.	2
T2	1	.	3	.	.
T3	1	2	3	.	2
T4	1	.	3	8	.
T5	1	6	3	8	6

Table 4D

Orbit length partitions of sets and sequences under G

G	2-sets	2-sequences
$G \leq A_4$		
T2	2^3	4^3
T4	6	12
$G \not\leq A_4$		
T1	2, 4	4^3
T3	2, 4	4, 8
T5	6	12

Table 5A: groups of degree 5

Group	Order	Even	Number of Classes	Name
T1	5	+	5	5
T2	10	+	4	D ₁₀
T3	20		5	5.4
T4	60	+	5	A ₅
T5	120		7	S ₅

Table 5B: group generators

$$a = (1, 2, 3, 4, 5) \quad c = (2, 3, 5, 4)$$

$$b = (1, 2)$$

$$\begin{array}{ll} T1 = \langle a \rangle & T4 = \langle a, bab \rangle \\ T2 = \langle a, c^2 \rangle & T5 = \langle a, b \rangle \\ T3 = \langle a, c \rangle & \end{array}$$

Table 5C: cycle type distribution

	1 ⁵	1 ³	2	2 ²	3	3 ²	4	5
T1	1	4	
T2	1	.	5	.	.	.	4	
T3	1	.	5	.	.	10	4	
T4	1	.	15	.	20	.	24	
T5	1	10	15	20	20	30	24	

Table 5D

Orbit length partitions of sets and sequences under G

G	2-sets	2-sequences
$G \triangle A_5$		
T1	5^2	5^4
T2	5^2	10^2
T4	10	20
$G \times A_5$		
T3	10	20
T5	10	20

Table 6A: groups of degree 6

Group	Order	Even	Imprimitive [2 ³] [3 ²]	Number of Classes	Other Representations	Name
T1	6		✓ ✓	6		6
T2	6		✓ ✓	3	3T2	Σ_3
T3	12		✓ ✓	6		D_{12}
T4	12	+	✓	4	4T4	A_4
T5	18		✓	9		$3 \times \Sigma_3$
T6	24		✓	8		$2 \times A_4$
T7	24	+	✓	5	4T5	$\Sigma_4/2^2$
T8	24		✓	5	4T5	$\Sigma_4/4$
T9	36		✓	9		$3^2 \cdot 2^2$
T10	36	+	✓	6		$3^2 \cdot 4$
T11	48		✓	10		$2 \times \Sigma_4$
T12	60	+		5	5T4	$L(2,5)$
T13	72		✓	9		$3^2 \cdot D_8$
T14	120			7	5T5	$PGL(2,5)$
T15	360	+		7		A_6
T16	720			11		Σ_6

Table 6B: group generators

$$\begin{array}{ll}
 a = (1,2,3) & i = (1,3)(2,4) \\
 b = (1,4)(2,5)(3,6) & j = (1,6)(2,5)(3,4) \\
 c = (1,5,2,4)(3,6) & k = (1,2,3,4,5) \\
 d = ab & l = (1,6)(2,5) \\
 e = bc^2 & m = (2,3,5,4) \\
 f = (1,2) & \\
 g = (1,3,5)(2,4,6) & \\
 h = fgf^{-2} &
 \end{array}$$

$$\begin{array}{ll}
 T_1 = \langle d \rangle & T_{11} = \langle f, g, i \rangle \\
 T_2 = \langle e, j \rangle & T_{12} = \langle k, l \rangle \\
 T_3 = \langle d, e \rangle & T_{13} = \langle a, b, c \rangle \\
 T_4 = \langle g, h \rangle & T_{14} = \langle k, l, m \rangle \\
 T_5 = \langle a, b \rangle & T_{15} = \langle c, k \rangle \\
 T_6 = \langle g, f \rangle & T_{16} = \langle d, k \rangle \\
 T_7 = \langle g, h, i \rangle & \\
 T_8 = \langle g, h, j \rangle & \\
 T_9 = \langle a, b, e \rangle & \\
 T_{10} = \langle a, c \rangle &
 \end{array}$$

Table 6C: cycle type distribution

	1^6	2^4	2^2	2^3	3^3	2^2	3^2	4^2	4^2	5	
T1	1	.	.	1	.	.	2	.	.	.	2
T2	1	.	.	3	.	.	2
T3	1	.	3	4	.	.	2	.	.	.	2
T4	1	.	3	.	.	.	8
T5	1	.	.	3	4	.	4	.	.	.	6
T6	1	3	3	1	.	.	8	.	.	.	8
T7	1	.	9	.	.	.	8	.	6	.	.
T8	1	.	3	6	.	.	8	6	.	.	.
T9	1	.	9	6	4	.	4	.	.	.	12
T10	1	.	9	.	4	.	4	.	18	.	.
T11	1	3	9	7	.	.	8	6	6	.	8
T12	1	.	15	.	.	.	20	.	.	24	.
T13	1	6	9	6	4	12	4	.	18	.	12
T14	1	.	15	10	.	.	20	30	.	24	20
T15	1	.	45	.	40	.	40	.	90	144	.
T16	1	15	45	15	40	120	40	90	90	144	120

Table 6D

Orbit length partitions of sets and sequences under G

G	2-sets	3-sets	2-sequences
<u>$G \leq A_6$</u>			
T4	3,12	$4^2, 6^2$	$6, 12^2$
T7	3,12	$4^2, 12$	$6, 24$
T10	6,9	2,18	12,18
T12	15	10^2	30
T15	15	20	30
<u>$G \times A_6$</u>			
T1	$3, 6^2$	$2, 6^3$	6^5
T2	$3^2, 6$	$2, 6^3$	6^5
T3	$3, 6^2$	$2, 6, 12$	$6, 12^2$
T5	6,9	2,18	$6^2, 18$
T6	3,12	$6^2, 8$	$6, 12^2$
T8	3,12	8,12	$6, 24$
T9	6,9	2,18	12,18
T11	3,12	8,12	$6, 24$
T13	6,9	2,18	12,18
T14	15	20	30
T16	15	20	30

Table 7A: groups of degree 7

Group	Order	Even	Number of Classes	Name
T1	7	+	7	7
T2	14		5	D ₁₄
T3	21	+	5	7·3
T4	42		7	7·6
T5	168	+	6	L(3,2)
T6	2520	+	9	A ₇
T7	5040		15	S ₇

Table 7B: group generators

$$a = (1,2,3,4,5,6,7)$$

$$c = (2,3)(4,7)$$

$$b = (2,4,3,7,5,6)$$

$$d = (1,2,3)$$

$$T1 = \langle a \rangle$$

$$T6 = \langle a, d \rangle$$

$$T2 = \langle a, b^3 \rangle$$

$$T7 = \langle b, d \rangle$$

$$T3 = \langle a, b^2 \rangle$$

$$T4 = \langle a, b \rangle$$

$$T5 = \langle a, c \rangle$$

Table-7C: cycle type distribution

	2	2 ²	2 ³	3	2	3	3 ²	4	2	4	5	5	6		
T1	17	15	13	1	14	12	22	1	13	1	3	12	2	1	7
T2	1	.	.	7	6	.	
T3	1	14	6	.	
T4	1	.	.	7	.	.	.	14	.	.	.	14	6	.	
T5	1	.	21	.	.	.	56	42	.	.	.	48	.	.	
T6	1	.	105	.	70	.	210	280	.	630	.	504	.	720	
T7	1	21	105	105	70	420	210	280	210	630	420	504	504	840	720

Table 7D

Orbit length partitions of sets and sequences under G

G	2-sets	3-sets	2-sequences
<u>$G \leq A_7$</u>			
T1	7^3	7^5	7^6
T3	21	$7^2, 21$	21^2
T5	21	$7, 28$	42
T6	21	35	42
<u>$G \times A_7$</u>			
T2	7^3	$7^3, 14$	14^3
T4	21	$14, 21$	42
T7	21	35	42

Table 8A: groups of degree 8

Group	Order	Even	Imprimitve [2 ⁴] [4 ²]	Number of Classes	Other Representations	Name
T1	8		/ /	8		8
T2	8	+	3 ✓	8		2x4
T3	8	+	7 ✓	8		2 ³
T4	8	+	5 ✓	5	4T3	D ₈
T5	8	+	/ ✓	5		Q ₈
T6	16		/ /	7		
T7	16		/ /	10		
T8	16		/ /	7		
T9	16	+	3 ✓	10		
T10	16	+	3 ✓	10		
T11	16	+	/ ✓	10		
T12	24	+	/			SL(2,3)
T13	24	+	/ ✓	8	6T6	2xA ₄
T14	24	+	/ ✓	5	4T5	Σ ₄
T15	32		/ ✓	11		
T16	32		/ ✓	11		
T17	32		/ ✓	14		
T18	32	+	/ ✓	14		
T19	32	+	/ ✓	11		
T20	32	+	/ ✓	11		
T21	32		/ ✓	11		
T22	32	+	/ ✓	17		
T23	48		/ ✓	8		
T24	48	+	/ ✓	10	6T11	2x Σ ₄

Table 8A (continued)

Group	Order	Even	Imprimitive [2 ⁴] [4 ²]	Number of Classes	Other Representation	Name
T25	56	+		8		3·7
T26	64		✓ ✓	16		
T27	64		✓ ✓	13		
T28	64		✓ ✓	13		
T29	64	+	✓ ✓	16		
T30	64		✓ ✓	13		
T31	64		✓ ✓	16		
T32	96	+	✓	11		
T33	96	+	✓	10		
T34	96	+	✓	10		
T35	128		✓ ✓	20		
T36	168	+		8		2 ³ ·(7·3)
T37	168	+		6	7T5	L(2,7)
T38	192		✓	16		
T39	192	+	✓	13		
T40	192		✓	13		
T41	192	+	✓	14		
T42	288	+	✓	14		
T43	336			9		PGL(2,7)
T44	384			20		
T45	576	+	✓	16		
T46	576		✓	13		
T47			✓	20		
T48	1344	+		11		2 ³ ·L(3,2)
T49	20160	+		13		A ₈
T50	40320	+		22		Σ ₈

Table 8B: group generators

$a = (1,4,6,8; 2,3,5,7)$	$q = (1,6,2,5)(3,7)(4,8)$
$b = (1,3,5,7)(2,4,6,8)$	$r = (5,6)$
$c = (1,6)(2,5)(3,8)(4,7)$	$s = (1,3)(2,4)$
$d = (1,8)(2,7)(3,6)(4,5)$	$t = (1,2)$
$e = (1,7)(2,8)(3,5)(4,6)$	$u = (1,5)(2,6)$
$f = (1,7)(2,8)(3,6)(4,5)$	$v = (3,4)$
$g = (1,7,2,8)(3,5,4,6)$	$w = (1,3)(2,4)(7,8)$
$h = (3,4)(7,8)$	$x = (2,4,3)(6,8,7)$
$i = (1,6)(2,5)(3,4)$	$y = (1,8)(2,5)(3,6)(4,7)$
$j = (1,6)(2,5)(3,7)(4,8)$	$z = (6,8,7)$
$k = (1,6)(2,5)$	$A = (1,2,3,4,5,6,7)$
$l = (1,3)(2,4)(5,8)(6,7)$	$B = (2,4,3,7,5,6)$
$m = (1,5)(2,6)(3,7)(4,8)$	$C = (2,3)(4,7)$
$n = (3,5,7)(4,6,8)$	$D = (1,8)(2,4)(3,7)(5,6)$
$o = (1,4)(2,3)(5,6)(7,8)$	$E = (1,8)(2,7)(3,4)(5,6)$
$p = (1,2)(7,8)$	$F = (1,7,3,5)(2,8,4,6)$

$T1 = \langle a \rangle$	$T26 = \langle a, f, b^2 \rangle$
$T2 = \langle b, c \rangle$	$T27 = \langle a, t \rangle$
$T3 = \langle b^2, e, c \rangle$	$T28 = \langle a, u \rangle$
$T4 = \langle b, d \rangle$	$T29 = \langle b, e, f \rangle$
$T5 = \langle a^2, g \rangle$	$T30 = \langle b, p, iku \rangle$
$T6 = \langle a, f \rangle$	$T31 = \langle q, e, t \rangle$
$T7 = \langle a, h \rangle$	$T32 = \langle e, j, n \rangle$
$T8 = \langle a, i \rangle$	$T33 = \langle F, x \rangle$
$T9 = \langle b, e, c \rangle$	$T34 = \langle vsy, x, y \rangle$

Table 8B (continued)

T10 = $\langle b, j \rangle$
 T11 = $\langle a^2, b^2, l \rangle$

T12 = $\langle g, n \rangle$

T13 = $\langle h, j, n \rangle$

T14 = $\langle n, o \rangle$

T15 = $\langle a, f, h \rangle$

T16 = $\langle a, b^2 \rangle$

T17 = $\langle a, e \rangle$

T18 = $\langle b, e, j \rangle$

T19 = $\langle b, f \rangle$

T20 = $\langle b, p \rangle$

T21 = $\langle q, e \rangle$

T22 = $\langle a^2, b^2, j, e \rangle$

T23 = $\langle n, w \rangle$

T24 = $\langle c, n, s \rangle$

T25 = $\langle A, D \rangle$

T35 = $\langle a, f, t \rangle$

T36 = $\langle A, D, B^2 \rangle$

T37 = $\langle A, B^2, E \rangle$

T38 = $\langle v, e, n \rangle$

T39 = $\langle j, n, s \rangle$

T40 = $\langle j, n, shv \rangle$

T41 = $\langle F, x, y \rangle$

T42 = $\langle s, z, m \rangle$

T43 = $\langle A, B, E \rangle$

T44 = $\langle t, b, s \rangle$

T45 = $\langle s, z, m, y \rangle$

T46 = $\langle s, z, q \rangle$

T47 = $\langle sv, t, m \rangle$

T48 = $\langle A, C, D \rangle$

T49 = $\langle A, z \rangle$

T50 = $\langle \text{?}, t \rangle$

Table 8C: cycle type distribution

	1^8	2	2^2	2^3	2^4	3	3	3^2	3^2	4	4	4
	1^6	1^4	1^2	1^3	1^5	1^3	1^1	1^2	1^2	1^4	1^2	1^2
T1	1	1
T2	1	3
T3	1	7
T4	1	5
T5	1	1
T6	1	.	.	.	4	5
T7	1	.	2	.	.	1
T8	1	.	.	4	1
T9	1	.	2	.	9
T10	1	.	2	.	5
T11	1	.	2	.	5
T12	1	.	.	.	1	.	.	.	8	.	.	.
T13	1	.	.	.	7	.	.	.	8	.	.	8
T14	1	.	.	.	9	.	.	.	8	.	.	.
T15	1	.	2	8	5
T16	1	.	6	.	5
T17	1	.	2	.	5	4	.	4
T18	1	.	6	.	13
T19	1	.	2	.	9	8	.
T20	1	.	6	.	5
T21	1	.	6	.	5	16
T22	1	.	6	.	13

Table 8C (continued)

	2	2^2	2^3	3	2	2^2	3^2	3^2	4	2	4		
	1^8	1^6	1^4	1^2	2^4	1^5	1^3	1	1^2	2	1^4	1^2	2^2
T23	1	.	.	12	1	.	.	8
T24	1	.	6	.	13	.	.	8
T25	1	.	.	.	7
T26	1	.	6	8	13	4	.	4	.
T27	1	4	6	4	5	8
T28	1	.	10	.	9	8	16	.
T29	1	.	10	.	17	8	.	.
T30	1	.	6	8	5	4	.	20	.
T31	1	4	6	4	13	24	.
T32	1	.	6	.	13	.	.	.	32
T33	1	.	6	.	13	.	.	.	32
T34	1	.	6	.	21	.	.	.	32
T35	1	4	10	12	17	4	8	28	.
T36	1	.	.	.	7	.	.	.	56
T37	1	.	.	.	21	.	.	.	56
T38	1	4	6	4	13	.	.	.	32	32	.	24	.
T39	1	.	18	.	25	.	.	.	32	.	.	24	.
T40	1	.	6	24	13	.	.	.	32	.	12	.	12
T41	1	.	18	.	25	.	.	.	32	.	.	24	.
T42	1	.	6	.	21	16	.	48	64
T43	1	.	.	28	21	.	.	.	56

Table 8C (continued)

						3	3					4	
	2	2^2	2^3		3	2	2^2	3^2	3^2	4	2	4	
	1^8	1^6	1^4	1^2	2^4	1^5	1^3	1	1^2	2	1^4	1^2	2^2
T44	1	4	18	28	25	.	.	.	32	32	12	24	36
T45	1	.	42	.	33	16	.	48	64	.	.	72	.
T46	1	.	42	.	9	16	.	48	64	.	.	72	144
T47	1	12	42	36	33	16	96	48	64	.	12	72	180
T48	1	.	42	.	49	.	.	.	224	.	.	168	.
T49	1	.	210	.	105	112	.	1680	1120	1	.	2520	.
T50	1	28	210	420	105	112	1120	1680	1120	1120	420	2520	1260

Table 8C (continued)

	4	5	2	5	6	6	7	
	3	5	1	3	1 ²	2	1	8
	1	4 ²	1 ³	1				
T1		2						4
T2		4						
T3								
T4		2						
T5		6						
T6		2						4
T7		4						8
T8		6						4
T9		4						
T10		8						
T11		8						
T12		6						8
T13								8
T14		6						
T15		8						8
T16		4						16
T17		8						8
T18		12						
T19		12						
T20		20						
T21		4						
T22		12						

Table 8C (continued)

	4	5	6	7	8
	3	2	1	2	1
	1	4 ²	1 ³	3	2
T23	6	.	.	8	12
T24	12	.	.	8	.
T25	48
T26	12	.	.	.	16
T27	20	.	.	.	16
T28	4	.	.	.	16
T29	28
T30	20
T31	12
T32	12	.	.	32	.
T33	12	.	.	32	.
T34	36
T35	28	.	.	.	16
T36	.	.	.	56	48
T37	42	.	.	.	48
T38	12	.	.	32	32
T39	60	.	.	32	.
T40	12	.	.	32	48
T41	60	.	.	32	.
T42	36	.	.	96	.
T43	42	.	.	56	48
T44	60	.	.	32	32
					48

Table 8C (continued)

	4	5	6	7	8
	3	2	5	6	7
	1	1 ²	1 ³	1 ²	1
T45	108			192	
T46	36				144
T47	96	108		192	144
T48		252		224	384
T49		1260	1344	2688	3360
T50	3360	1260	1344	4032	2688
				3360	5760
					5040

Table 8D

Orbit length partitions of sets and sequences under G

G	2-sets	3-sets	4-sets	2-sequences
<u>$G < A_8$</u>				
T2	$4^3, 8^2$	8^7	$2^3, 4^2, 8^7$	8^7
T3	4^7	8^7	$2^7, 8^7$	8^7
T4	$4^5, 8$	8^7	$2^3, 4^4, 8^6$	8^7
T5	$4, 8^3$	8^7	$2^3, 8^8$	8^7
T9	$4^3, 8^2$	$8^3, 16^2$	$2^3, 4^2, 8^3, 16^2$	$8^3, 16^2$
T10	$4^3, 16$	$8^3, 16^2$	$2, 4^3, 8^3, 16^2$	$8^3, 16^2$
T11	$4, 8^3$	$8^3, 16^2$	$2^3, 8^4, 16^2$	$8^3, 16^2$
T12	$4, 24$	$8, 24^2$	$6, 8^2, 24^2$	$8, 24^2$
T13	$4, 12^2$	$8, 24^2$	$2, 6^2, 8, 24^2$	$8, 24^2$
T14	$4, 12^2$	$8, 24^2$	$2, 6^2, 8, 12^2, 24$	$8, 24^2$
T18	$4^3, 16$	$8, 16^3$	$2, 4^3, 8^3, 32$	$8^3, 32$
T19	$4, 8, 16$	$8, 16, 32$	$2, 4, 8^2, 16, 32$	$8, 16, 32$
T20	$4, 8, 16$	$8^3, 32$	$2, 4, 8^2, 16^3$	$8, 16^3$
T22	$4, 8^3$	$8^3, 32$	$2^3, 8^2, 16^3$	$8, 16^3$
T24	$4, 12^2$	$8, 24^2$	$2, 6^2, 8, 24^2$	$8, 24^2$
T25	28	56	14, 56	56
T29	$4, 8, 16$	$8, 16, 32$	$2, 4, 8^2, 16, 32$	$8, 16, 32$
T32	$4, 24$	$24, 32$	$6, 8^2, 48$	$8, 48$
T33	$12, 16$	$8, 48$	$2, 12, 24, 32$	$24, 32$
T34	$12, 16$	$8, 48$	$2, 12^3, 32$	$24, 32$
T36	28	56	14, 56	56
T37	28	56	$14^2, 42$	56

Table 8D (continued)

G <u>G < A</u> 8	2-sets	3-sets	4-sets	2-sequences
T39	4,24	24,32	6,8 ² ,48	8,48
T41	12,16	8,48	2,12,24,32	24,32
T42	12,16	8,48	2,32,36	24,32
T45	12,16	8,48	2,32,36	24,32
T48	28	56	14,56	56
T49	28	56	70	56

Table 8D (continued)

G	2-sets	3-sets	4-sets	2-sequences
<u>GKA₈</u>				
T1	4,8 ³	8 ⁷	2,4,8 ⁸	8 ⁷
T6	4,8 ³	8 ³ ,16 ²	2,4,8 ⁴ ,16 ²	8,16 ³
T7	4,8,16	8 ³ ,16 ²	2,4,8 ² ,16 ³	8 ³ ,16 ²
T8	4,8,16	8 ³ ,16 ²	2,4,8 ² ,16 ³	8,16 ³
T15	4,8,16	8,16 ³	2,4,8 ² ,16,32	8,16,32
T16	4,8,16	8 ³ ,32	2,4,16 ⁴	8,16 ³
T17	4,8,16	8,16,32	2,4,16 ² ,32	8 ³ ,32
T21	4,8 ³	8 ³ ,32	2 ³ ,16 ⁴	8,16 ³
T23	4,24	8,24 ²	6,16,24 ²	8,48
T26	4,8,16	8,16,32	2,4,16 ² ,32	8,16,32
T27	4,8,16	8 ³ ,32	2,4,16 ⁴	8,16 ³
T28	4,8,16	8,16,32	2,4,16 ² ,32	8,16,32
T30	4,8,16	8,16,32	2,4,16 ² ,32	8,16,32
T31	4,8 ³	8 ³ ,32	2 ³ ,16 ⁴	8,16 ³
T35	4,8,16	8,16,32	2,4,16 ² ,32	8,16,32
T38	4,24	24,32	6,16,48	8,48
T40	4,24	24,32	6,16,48	8,48
T43	28	56	28,42	56
T44	4,24	24,32	6,16,48	8,48
T46	12,16	8,48	2,32,36	24,32
T47	12,16	8,48	2,32,36	24,32
T50	28	56	70	56

APPENDIX 2

POLYNOMIALS WITH GIVEN TRANSITIVE GALOIS GROUPS OVER Q OF DEGREE UP TO 7

It is an unsolved problem whether any permutation group can appear as the Galois group of a polynomial over \mathbb{Q} . For each solvable group G it is known that there exists a polynomial $f(x)$ in $\mathbb{Q}[x]$ such that $\text{Gal}(f/\mathbb{Q}) = G$ (see [SHA]); however there does not appear to be a published general method of constructing this $f(x)$ given any solvable G .

For each transitive permutation group G of degree 3 to 7, we have computed a polynomial $f(x)$ such that $\text{Gal}(f/\mathbb{Q}) = G$. These polynomials appear in Table A2.1. The notation nTi means group T_i of degree n . The splitting field of $f(x)$ over \mathbb{Q} is denoted by $\text{spl}(f)$ and z_n denotes a primitive n -th root of 1.

For each polynomial $f(x)$ in Table A2.1, we proved that $\text{Gal}(f/\mathbb{Q})$ is the group indicated. Many of the polynomials $f(x)$ are constructed so that $\text{spl}(f)$ is contained in some known field. The methods of doing this include constructing $f(x)$ to be a resolvent polynomial, constructing $f(x)$ to be a composite polynomial, or if $\text{Gal}(f/\mathbb{Q})$ is to be cyclic, by constructing $f(x)$ such that $\text{spl}(f)$ is contained in $\mathbb{Q}(z_p)$, p prime (see [VDW, p. 163-168]). This knowledge about $\text{spl}(f)$ is

used to reduce or eliminate the work necessary to determine $\text{Gal}(f/Q)$. In fact, the only polynomials whose Galois groups are determined using other information than the splitting field, cycle types or discriminant are those $f(x)$ with $\text{Gal}(f/Q) = 5T2, 7T2, 7T3,$ or $7T5$. These exceptions are proved to have the group indicated by using the factorization of specific linear resolvents.

Given G , to find monic $f(x)$ in $\mathbb{Z}[x]$ such that $\text{Gal}(f/Q) = G$, where it is non-trivial to construct an appropriate splitting field, we do computer searching. If it is required that $\text{disc}(f)$ is a square we proceed in the following way:

Let p be an odd prime. $\text{disc}(f)$ a square implies $\text{disc}(f)$ is a quadratic residue (square) mod p or $\text{disc}(f) \pmod p = 0$. ($(p-1)/2$ elements in \mathbb{Z}_p are quadratic residues.) Note that $\text{disc}(f) \pmod p = \text{disc}(f \pmod p)$, where $\text{disc}(f \pmod p)$ is calculated over \mathbb{Z}_p (see Section 4.1). We can calculate $\text{disc}(f \pmod p)$ efficiently using equation (1.1), and for non-zero d in \mathbb{Z}_p , we use Euler's criterion [LON,p.111] that d is a quadratic residue if and only if $d^{(p-1)/2} = 1$.

We search over all polynomials in a given set, and return those $f(x)$ such that $\text{disc}(f \pmod {p_i}) = 0$ or $\text{disc}(f \pmod {p_i})$ is a quadratic residue, for all (small, consecutive) odd primes p_i in a given set S . In practice $30 \leq |S| \leq 40$. The discriminants over \mathbb{Z} of the found

polynomials $f(x)$ are then calculated as well as $\text{Gal}(f/\mathbb{Q})$.

The above search method to find polynomials with square discriminants is faster than a method which forms the complete discriminant over \mathbb{Z} for every polynomial $f(x)$ tested, because $f(x)$ can be rejected as soon as $\text{disc}(f \bmod p)$ is found to be a non-residue for some odd prime p . Also, working mod p allows us to perform this search using a very limited integer size. For example, this search has been implemented on the PDP-11/34 in the language PASCAL; using only 16 bits to represent an integer.

One can also search for monic $f(x)$ in $\mathbb{Z}[x]$ such that $\text{disc}(f \bmod p_i) = 0$ or the factor type of $f(x) \bmod p_i$ is the cycle type of some permutation in the required group of $f(x)$ (for all (small, consecutive) primes p_i in a fixed set).

McKay and Rohlicek use this technique, and in this way they found the polynomial in Table A2.1 with Galois group 7T2.

Table A2.1
Polynomials $f(x)$ such that $\text{Gal}(f/\mathbb{Q}) = G$.

G	disc(f)	$f(x)$	Remarks
Degree 3			
T1	7^2	x^3+x^2-2x-1	$\text{spl}(f)=\mathbb{Q}(z_7+z_7^{-1})$
T2	$-2^2 3^3$	x^3+2	
Degree 4			
T1	5^3	$x^4+x^3+x^2+x+1$	$\text{spl}(f)=\mathbb{Q}(z_5)$
T2	2^8	x^4+1	$\text{spl}(f)=\mathbb{Q}(z_8)$
T3	-2^{11}	x^4-2	
T4	$2^{12} 3^4$	$x^4+8x+12$	
T5	229	x^4+x+1	
Degree 5			
T1	11^4	$x^5+x^4-4x^3-3x^2+3x+1$	$\text{spl}(f)=\mathbb{Q}(z_{11}+z_{11}^{-1})$
T2	$2^{12} 5^6$	$x^5-5x+12$	
T3	$2^4 5^5$	x^5+2	
T4	$2^{16} 5^6$	$x^5+20x+16$	
T5	19.151	x^5-x+1	

Table A2.1 (continued)

G	disc(f)	f(x)	Remarks
Degree 6			
T1	-7 ⁵	$x^6 + x^5 + x^4 + x^3 + x^2 + x + 1$	$\text{spl}(f) = Q(z_7)$
T2	-2 ¹⁶ 3 ²¹	$x^6 + 108$	$\text{spl}(f) = \text{spl}(x^3 + 2)$
T3	-2 ¹¹ 3 ⁶	$x^6 + 2$	
T4	2 ⁶ 3 ⁸	$x^6 - 3x^2 - 1$	$\text{spl}(f) = \text{spl}(x^4 + 8x + 12)$
T5	-3 ¹¹	$x^6 + x^3 + 3$	
T6	-2 ⁶ 3 ⁸	$x^6 - 3x^2 + 1$	$\text{Gal}(x^3 - 3x + 1/Q) \cong A_3$
T7	2 ⁶ 229 ²	$x^6 - 4x^2 - 1$	$\text{spl}(f) = \text{spl}(x^4 + x + 1)$
T8	229 ³	$x^6 - 3x^5 + 6x^4 - 7x^3 + 2x^2 + x - 4$	$\text{spl}(f) = \text{spl}(x^4 + x + 1)$
T9	2 ⁸ 3 ⁹	$x^6 + 2x^3 - 2$	
T10	2 ¹⁰ 3 ⁶ 5 ⁴	$x^6 + 6x^4 + 2x^3 + 9x^2 + 6x - 4$	$f(x) = (x^3 + 3x + 1)^2 - 5$
T11	-2 ¹¹ 5 ² 7 ²	$x^6 + 2x^2 + 2$	
T12	2 ³⁶ 5 ⁸	$x^6 + 10x^5 + 55x^4 + 140x^3 + 175x^2 + 170x + 25$	$\text{spl}(f) = \text{spl}(x^5 + 20x + 16)$
T13	-2 ⁸ 733	$x^6 + 2x^4 + 2x^3 + x^2 + 2x + 2$	$f(x) = (x^3 + x + 1)^2 + 1$
T14	5 ²⁰ 19 ³ 151 ³	$x^6 + 10x^5 + 55x^4 + 140x^3 + 175x^2 - 3019x + 25$	$\text{spl}(f) = \text{spl}(x^5 - x + 1)$
T15	2 ¹⁶ 3 ⁶ 5 ⁶	$x^6 + 24x - 20$	
T16	-2 ⁶ 89.227	$x^6 + 2x + 2$	
Degree 7			
T1	29 ⁶ 17 ²	$x^7 + x^6 - 12x^5 - 7x^4 + 28x^3 + 14x^2 - 9x + 1$	$\text{spl}(f) = Q(z_{29} + z_{29}^{12} + z_{29}^{-1} + z_{29}^{-12})$
T2	-3 ⁶ 7 ⁹	$x^7 + 7x^3 + 7x^2 + 7x - 1$	
T3	2 ⁶ 7 ¹⁰	$x^7 - 14x^5 + 56x^3 - 56x + 22$	
T4	-2 ⁶ 7 ⁷	$x^7 + 2$	
T5	7 ⁸ 17 ²	$x^7 - 7x^3 + 14x^2 - 7x + 1$	
T6	3 ⁶ 7 ⁸	$x^7 + 7x^4 + 14x + 3$	
T7	-2 ⁶ 5.233.787	$x^7 + 2x + 2$	

APPENDIX 3

POLYNOMIALS WITH PSL(3,2) AS GALOIS GROUP OVER Q

Polynomials $f(x)$ such that $\text{Gal}(f/Q) = \text{PSL}(3,2)$ (group T5 in Table 7A of Appendix 1) have attracted interest over many years ([ERB, LAM, TRI] and their references). Recently, LaMacchia [LAM] has constructed an infinite family of polynomials with $\text{PSL}(3,2)$ as Galois group over Q .

In Table A3.1 we list integral $f(x)$ such that $\text{Gal}(f/Q) = \text{PSL}(3,2)$. We found these $f(x)$ by searching for polynomials with square discriminants as described in Appendix 2. All polynomials searched are of the form:

$$f(x) = x^7 + \sum_{i=0}^5 a_i x^i,$$

where $7|a_i$ and $|a_i| \leq M$ for $i=1, \dots, 5$; and $1 \leq a_0 \leq 2M$. The sets of polynomials searched are:

- (1) $\{f(x) : M = 14\}$,
- (2) $\{f(x) : M = 28, \text{ and } a_4, a_2 = 0\}$,
- (3) $\{f(x) : M = 56, a_4, a_2 \neq 0, \text{ and } a_5, a_3, a_1 \text{ are of the form } 7(2^i)\}$.

We know of no monic $f(x)$ in $Z[x]$ such that $\text{Gal}(f/Q) = \text{PSL}(3,2)$ and $\text{disc}(f) < 7^8 17^2$.

We did some similar searching in an effort to find degree 11 polynomials $f(x)$ such that $\text{Gal}(f/Q) = \text{PSL}(2,11)$ or

$\text{Gal}(f/\mathbb{Q}) = M_{11}$, the sporadic simple group of Mathieu of order 7920. No such $f(x)$ was found.

Table A3.1

Polyhomials $f(x)$ such that $\text{Gal}(f/\mathbb{Q}) = \text{PSL}(3,2)$

disc(f)	$f(x)$
$2^6 3^6 5^4 7^8$	$x^7 - 14x^5 + 14x^4 + 14x^3 - 14x^2 + 2$
$3^6 7^8 11^2 73^2$	$x^7 - 14x^5 - 7x^4 + 7x^3 - 7x^2 + 11$
$2^{10} 7^8$	$x^7 - 7x^5 - 14x^4 - 7x^3 - 7x + 2$
$7^8 17^2$	$x^7 - 7x^5 - 7x^4 + 7x^3 + 14x^2 + 7x + 2$
$2^6 3^4 7^8$	$x^7 - 7x^5 - 7x^4 + 7x^3 + 14x^2 + 7x + 3$
$2^6 7^8 13^2 19^2$	$x^7 - 7x^5 - 7x^4 + 14x^3 + 14x^2 - 14x + 6$
$2^{10} 5^2 7^8$	$x^7 - 7x^5 + 7x^3 - 7x + 4$
$2^{12} 7^8 23^2$	$x^7 - 7x^5 + 7x^3 + 14x^2 - 14x + 8$
$2^{14} 7^8$	$x^7 - 7x^5 + 14x^3 - 14x + 8$
$2^{12} 7^8 17^2$	$x^7 - 7x^5 + 7x^4 - 7x^3 + 7x^2 + 7$
$2^6 7^{10} 23^2$	$x^7 - 7x^5 + 7x^4 + 7x^3 - 14x^2 + 7x + 13$
$2^6 5^6 7^8$	$x^7 - 7x^5 + 7x^4 + 14x^3 - 14x^2 - 14x + 6$
$2^6 7^8 17^4$	$x^7 - 7x^5 + 7x^4 + 14x^3 - 14x^2 + 14x + 2$
$2^6 5^6 7^8 29^2$	$x^7 - 14x^4 - 7x^3 + 14x^2 + 14$
$2^{16} 5^2 7^8$	$x^7 - 14x^4 + 14x^3 + 4$
$2^8 7^8 643^2$	$x^7 - 14x^3 - 14x^2 + 7x + 22$
$7^8 17^2$	$x^7 - 7x^3 + 14x^2 - 7x + 1$
$3^8 7^8$	$x^7 - 7x + 3$ (example of Trinks [TRI])

Table A3.1 (continued)

disc(f)	f(x)
$5^2 7^8 13^2 17^2$	$x^7 + 7x^4 - 7x^3 - 7x^2 + 14x + 3$
$3^6 5^2 7^8 13^2 17^2$	$x^7 + 14x^4 - 7x^3 - 14x^2 - 14x + 13$
$3^4 7^8 13^2 19^2 41^2$	$x^7 + 14x^5 - 7x^4 - 7x^3 - 7x^2 - 14x + 17$
$2^{12} 5^2 7^8 149^2$	$x^7 - 28x^3 + 28x + 20$
$3^{12} 7^8 47^2$	$x^7 - 21x^3 + 7x + 27$
$2^{20} 7^8 457^2$	$x^7 - 14x^5 - 28x^3 + 28x + 16$
$2^{12} 7^8 11^2 2699^2$	$x^7 - 56x^3 + 28x + 44$