

Technical Report No. 1/20, March 2020

A NUMERICAL STUDY OF ENTROPY AND RESIDUAL
ENTROPY ESTIMATORS BASED ON SMOOTH DENSITY
ESTIMATORS FOR NON-NEGATIVE RANDOM VARIABLES

Yogendra P. Chaubey and Nhat Linh Vu

A Numerical Study of Entropy and Residual Entropy Estimators Based on Smooth Density Estimators for Non-negative Random Variables

YOGENDRA P. CHAUBEY

Department of Mathematics and Statistics
Concordia University, Montreal, Canada H4B 1R6
E-mail: yogen.chaubey@concordia.ca

NHAT LINH VU

Department of Mathematics and Statistics
Concordia University, Montreal, Canada H4B 1R6
E-mail: nhatlinh.vu@concordia.ca

Abstract

In this paper, we are interested in the entropy of a non-negative random variable. Since the underlying probability density function is unknown, we propose the use of Poisson smoothed histogram density estimator in order to estimate the entropy. To study the performance of our estimator, we run simulations on a wide range of densities and compare our entropy estimators with the existing estimators that based on different approaches such as spacing estimators. Furthermore, we extend our study to residual entropy estimators which is the entropy of a random variable given that it has been survived up to time t .

Keywords: entropy estimator, information theory, residual entropy, density estimator, survival function.

1 Introduction

For a continuous random variable X , the differential entropy $H(X)$ is defined as

$$H(X) = - \int_{-\infty}^{\infty} f(x) \log f(x) dx. \quad (1)$$

In the literature, it is also often referred as Shannon entropy as it generalizes the concept of entropy for a discrete random variable proposed by by Shanon (1948) in his landmark paper, that is given by

$$H(X) = - \sum_{x \in A} \mathbb{P}(x) \log \mathbb{P}(x), \quad (2)$$

where A is the set of possible values of X , and $\mathbb{P}(X)$ denotes its probability mass function (*pmf*). We will simply refer to $H(X)$ as the entropy of the random variable X .

The concept of entropy takes a central place in statistical theory and applications (see Cover and Thomas, 1991 Kapur, 1993 and Kapur and Kesavan, 1992). One of the well known applications of entropy in statistics is the test of normality for a random variable because of the characterizing property that the normal distribution attains the maximum entropy among all continuous distributions with a given variance [see Vacicek, 1976 and an adaptation to testing exponentiality by Chaubey, Mudholkar and Smethurst, 1993].

Ebrahimi (1996) provided an interpretation of $H(X)$ as a measure of uncertainty associated with f . In the context of life testing, X being the lifetime of a unit, knowing that the unit has survived up to time t , a more appropriate measure for uncertainty in f is defined as

$$H(X, t) = - \int_t \left(\frac{f(x)}{R(t)} \right) \left(\log \frac{f(x)}{R(t)} \right), \quad (3)$$

where $R(t) = \mathbb{P}(X > t)$ is the reliability or the survival function. $H(X, t)$ is called *residual entropy* of X given the event $\{X > t\}$. In this paper, we focus on estimating $H(X)$ and more generally $H(X, t)$ when X is continuous and more specifically, when X is a non-negative. In recent decades, there exist a huge literature on entropy estimators, which can be classified as follows.

- (i) A naive approach is to discretize the support of the underlying density function into m bins, then for each bin we compute the empirical measures, corresponding to $p_i = \int_{A_i} f(x)dx$. That is if X_1, X_2, \dots, X_n are independently and identically distributed (i.i.d.) continuous random variables, then $\hat{p}_i = \frac{1}{n} \sum_{j=1}^n \mathbf{1}_{A_i}(X_j)$, where $\{A_i\}_{i=1}^m$ are mutually disjoint subsets of \mathbb{R} such that $\cup_{i=1}^m A_i = \text{support}(X)$. Clearly, \hat{p}_i is the maximum likelihood estimator (MLE) for p_i . With this estimator \hat{p}_i , a “naive” or “plug-in” MLE estimator of entropy is given as

$$\hat{H}^{MLE}(X) = - \sum_{i=1}^m \hat{p}_i \log \hat{p}_i. \quad (4)$$

It has been shown that this MLE estimator of entropy results in heavy bias. Consequently, the problem of bias correction and the choice of m have drawn the attention of many researchers; see Paninski (2003) for a detailed account.

- (ii) One straightforward approach for entropy estimation is to estimate the underlying density function $f(x)$ by some well-known density estimator $\hat{f}_n(x)$, then plug it into (??) to obtain the entropy estimator

$$\hat{H}^{PlugIn}(X) = - \int_0^\infty \hat{f}_n(x) \log \hat{f}_n(x) dx. \quad (5)$$

The *fixed symmetric kernel density estimator* $\hat{f}_n^{Fixed}(x)$, which is already a well-known and popular approach for estimating the pdf with an unbounded support, is an example of density estimator which is defined as

$$\hat{f}_n^{Fixed}(x) = \frac{1}{nb} \sum_{i=1}^n K \left(\frac{x - X_i}{b} \right), \quad (6)$$

where K is a symmetric density function with mean zero and variance one, and b is the smoothing parameter, called the bandwidth. It is obvious that the performance of this plugin entropy estimator totally depends on the density estimator \hat{f}_n . However, when dealing with non-negative random

variables, this $\hat{f}_n^{Fixed}(\cdot)$ is shown to produce a heavy bias near the boundary. Consequently, it would result in a better estimator if we replace \hat{f}_n^{Fixed} by some density estimators that is free of boundary effect.

- (iii) Motivated by the representation of entropy as an expected value

$$H(X) = - \int_0^\infty f(x) \log f(x) dx = -\mathbb{E}[\log f(X)].$$

By the strong law of large number we have $-\frac{1}{n} \sum_{i=1}^n \log f(X_i) \xrightarrow{a.s.} H(X)$. Thus we obtain a new entropy estimator if we replace $f(\cdot)$ by an appropriate density estimator $\hat{f}_n(\cdot)$, as given by

$$\hat{H}^{Meanlog}(X) = -\frac{1}{n} \sum_{i=1}^n \log \hat{f}_n(X_i). \quad (7)$$

- (iv) The entropy can be estimated by another approach, called “spacing”, which is initiated by Vasicek (1975). By the change of variable $p = F(x)$, the entropy (??) can be expressed in the form

$$H(X) = - \int_{-\infty}^\infty f(x) \log f(x) dx = \int_0^1 \log \left\{ \frac{d}{dp} F^{-1}(p) \right\} dp.$$

To estimate $H(X)$, the distribution F is replaced by the empirical distribution F_n , and the differential operator is replaced by the difference operator. As a result, the derivative of $F^{-1}(p)$ is estimated by $\frac{n}{2m}(X_{(i+m)} - X_{(i-m)})$ for $(i-1)/n < p \leq i/n$, $i = m+1, m+2, \dots, n-m$, where $X_{(i)}$'s are the order statistics and m is a positive integer smaller than $n/2$. When $p \leq m/n$ or $p > (n-m)/n$, one-sided differences are used. That is, $(X_{(i+m)} - X_{(1)})$ and $(X_{(n)} - X_{(i-m)})$ are in place of $(X_{(i+m)} - X_{(i-m)})$ respectively. All together this leads to the following estimator of entropy

$$\hat{H}^{Vasicek}(X) = \frac{1}{n} \sum_{i=1}^n \log \left\{ \frac{n}{2m}(X_{(i+m)} - X_{(i-m)}) \right\}. \quad (8)$$

Motivated by the idea of spacing, researchers have followed this direction and proposed other versions of entropy estimator, which are claimed to have a better performance. We will list some of them in the next section. One of the greatest weakness of this spacing estimator is the choice of spacing parameter m , which does not have the optimal form.

- (v) Lastly, different from all above estimators, Bouzebda et al. (2013) presented a potentially estimator of entropy based on smooth estimator of quantile density function. Their idea again starts with the expression of the entropy

$$H(X) = \int_0^1 \log \left\{ \frac{d}{dp} F^{-1}(p) \right\} dp = \int_0^1 \log \left\{ \frac{d}{dp} Q(p) \right\} dp = \int_0^1 \log q(p) dp,$$

where $Q(p) := \inf\{t : F(t) \geq p\}$ for $0 \leq p \leq 1$ is the quantile function and $q(p) := dQ(p)/dp = 1/f(Q(p))$ is the quantile density function. Then a new entropy estimator can be obtained by substituting $q(\cdot)$ by its appropriate estimator $\hat{q}_n(\cdot)$. That is

$$\hat{H}^{Quantile}(X) = \int_0^1 \log \hat{q}_n(p) dp. \quad (9)$$

Bouzebda et al. (2013) were motivated by the work of Cheng and Parzen (1997), which introduced a kernel type estimator $\hat{q}_n(\cdot)$ that has good asymptotic properties.

For estimating the residual entropy $H(X, t)$ we can write (??) as suggested in Belzunce et al. (2001)

$$H(X, t) = \log(R(t)) - \frac{1}{R(t)} \int_t^\infty f(x) \log f(x) dx. \quad (10)$$

Then the residual entropy can be estimated if we replace $R(t)$ by an empirical or kernel estimator $\hat{R}(t)$, and an estimator $\hat{g}_n(x) = \hat{f}_n(x) \log \hat{f}_n(x)$ in place of the functional $g(x) = f(x) \log f(x)$.

The organization of the paper is as follows. We propose our entropy and residual entropy estimators corresponding to nonnegative random variables in Section 2. The first part of Section 3 is dedicated to the entropy estimators comparison, in which we compare our entropy estimators to the existing entropy estimators by running simulations on a wide range of densities. Finally, we perform the comparison of our residual entropy with other estimators in the second part of Section 3.

2 Main results

2.1 Entropy estimators

In this sub-section, we will propose two entropy estimators based on the idea of (??) and (??). The asymptotic properties of these estimators, such as asymptotic convergence and asymptotic normality in the context of densities with non-negative support can be established along the same lines as in Hall and Morton (1993) for estimators of type (??) and Eggermont and LaRiccia (1999) for estimators of type (??). We defer the discussion of these in another publication.

Recall that the entropy of a random variable can be estimated by the direct plugin approach:

$$\hat{H}^{Plugin}(X) = - \int_0^\infty \hat{f}_n(x) \log \hat{f}_n(x) dx,$$

or by the sample mean of $\log f(X_i), i = 1, 2, \dots, n$:

$$\hat{H}^{Meanlog}(X) = -\frac{1}{n} \sum_{i=1}^n \log \hat{f}_n(X_i).$$

We observe that both the entropy estimators presented above mainly depend on the density estimator, so we would expect to obtain a good entropy estimator if the chosen density estimator is well-behaved. In literature, regarding to non-negative random variable, there are so many candidates for the density estimators with nice asymptotic properties. Among these, we especially focus on the Poisson smoothed histogram density estimator, which has the following form (see Chaubey and Sen, 2009),

$$\hat{f}^{Pois}(x) = k \sum_{i=0}^{\infty} \left[F_n\left(\frac{i+1}{k}\right) - F_n\left(\frac{i}{k}\right) \right] e^{-kx} \frac{(kx)^i}{i!}, \quad (11)$$

where $F_n(\cdot)$ is the empirical distribution function, and $k = k(n)$ can be viewed as the smoothing parameter. This estimator $\hat{f}^{Pois}(x)$ can be interpreted as a random weighted sum of Poisson mass functions.

The asymptotic properties of $\hat{f}^{Pois}(x)$ have been studied and its first weak convergence was proven by Bouezmarni and Scaillet (2005) under the assumption $\lim_{n \rightarrow \infty} k_n = \infty$ and $\lim_{n \rightarrow \infty} nk_n^{-2} = \infty$. They also obtained the weak convergence for the case of unbounded pdf f at $x = 0$. Later on, Chaubey et al. (2010) filled the gap in the asymptotic theory of \hat{f}^{Pois} by proposing the asymptotic bias, asymptotic variance, strong consistency and the asymptotic normality of the estimator. Particularly, under the assumptions that $k_n = cn^h$ for some constant c and $0 < h < 1$, and $f'(x)$ satisfies the Lipschitz condition of order $\alpha > 0$. The asymptotic bias and variance of $\hat{f}^{Pois}(\cdot)$ are given by

$$Bias[\hat{f}^{Pois}(x)] \approx \frac{f'(x)}{2cn^h}, \quad (12)$$

$$Var[\hat{f}^{Pois}(x)] \approx \frac{\mathbb{E}[X]}{2} \sqrt{\frac{c}{2\pi x^3}} f(x) n^{h/2-1}. \quad (13)$$

For the strong consistency of $\hat{f}^{Pois}(x)$, if $\mathbb{E}[X^{-2}] < \infty$, $f'(x)$ is bounded and $k_n = O(n^h)$, then

$$\|\hat{f}^{Pois}(x) - f(x)\| \xrightarrow{a.s.} 0.$$

For the asymptotic normality of $\hat{f}^{Pois}(x)$, if $\mathbb{E}[X^{-2}] < \infty$, $k_n = O(n^{2/5})$, and $f'(x)$ satisfies the Lipschitz order α condition, then for x in a compact set $I \subset \mathbb{R}^+$,

$$n^{2/5}(\hat{f}^{Pois}(x) - f(x)) - \frac{1}{2\delta^2} f'(x) \xrightarrow{D} \mathcal{G},$$

where \mathcal{G} is the Gaussian process with covariance function $\gamma_x^2 \delta_{x,s}$, where $\gamma_x^2 = \frac{\mathbb{E}[X]}{2} (2\pi x^3)^{-1/2} f(x) \delta$, $\delta_{x,s} = 0$ for $x \neq s$, $\delta_{x,s} = 1$ for $x = s$, and $\delta = \lim_{n \rightarrow \infty} (n^{-1/5} k_n^{1/2})$.

Similar to the problem of bandwidth selection in kernel density estimators, the choice of the smoothing parameter k strongly affects the performance of the resulting density estimators. In general, large values of k correspond to over-smoothing, whereas small values of k correspond to under-smoothing.

With these nice asymptotic properties of $\hat{f}^{Pois}(\cdot)$, we propose here two entropy estimators of the form

$$\hat{H}^{Plugin-Pois}(X) = - \int_0^\infty \hat{f}^{Pois}(x) \log \hat{f}^{Pois}(x) dx, \quad (14)$$

$$\hat{H}^{Meanlog-Pois}(X) = - \sum_{i=1}^n \log \hat{f}^{Pois}(X_i). \quad (15)$$

2.2 Residual entropy estimator

Motivated by the well-behaviour of $\hat{f}^{Pois}(\cdot)$, we suggest a direct plugin residual entropy estimator where the $\hat{f}^{Pois}(x)$ is in place of $f(x)$. That is, our proposed residual entropy estimator is of the form

$$\hat{H}^{Plugin-Pois}(X, t) = \log(\hat{R}(t)) - \frac{1}{\hat{R}(t)} \int_t^\infty \hat{f}^{Pois}(x) \log \hat{f}^{Pois}(x) dx, \quad (16)$$

where $\hat{R}(t) = \int_t^\infty \hat{f}^{Pois}(x) dx$.

3 Simulation studies and discussion

To study the performance of entropy and residual entropy estimators, we run simulations for a wide range of densities which consists fifteen non-negative densities below. They are categorized into three groups: monotone density, unimodal density, and bimodal density.

1. Monotone density

- Standard Exponential(1) with true $H(X) = 1$.
- Exponential(10) with true $H(X) = -\log 10 - 1 \approx -1.3026$.
- Pareto(2,1) with true $H(X) \approx 0.8069$.
- Log-Normal(0,2) with true $H(X) \approx 2.1121$.
- Weibull(0.5,0.5) with true $H(X) \approx 0.4228$.

2. Unimodal density

- Gamma(2,2) with true $H(X) \approx 0.8841$.
- Gamma(7.5,1) with true $H(X) \approx 2.3804$.
- Log-Normal(0,0.5) with true $H(X) \approx 0.7258$.
- Maxwell(1) with true $H(X) \approx 0.9962$.
- Maxwell(20) with true $H(X) \approx -0.5017$.
- Weibull(2,2) with true $H(X) \approx 1.2886$.

3. Bimodal density

- Mix Gamma: $(1/2)\text{Gamma}(0.5,0.5)+(1/2)\text{Gamma}(2,2)$ with true $H(X) \approx 2.2757$.
- Mix Lognorm: $(1/2)\text{Lognorm}(0,0.5)+(1/2)\text{Lognorm}(0,2)$ with true $H(X) \approx 1.6724$.
- Mix Maxwell: $(1/2)\text{Maxwell}(1)+(1/2)\text{Maxwell}(20)$ with true $H(X) \approx 0.8014$.
- Mix Weibull: $(1/2)\text{Weibull}(0.5,0.5)+(1/2)\text{Weibull}(2,2)$ with true $H(X) \approx 1.1330$.

The densities in the monotone group are chosen with different rates of decay, from the slowest standard Exp(1) to the fastest Exp(10). The Pareto distribution is included here to show the effect of different support $[1, \infty)$ on entropy estimation. On the contrary to the first group, the second group consists of well-behaved densities from highly concentrated Maxwell(20) to widely spread Gamma(7.5,1). Lastly, the simulations are extended to the bimodal densities where each density is a mixture of the same family density but with different parameters. Note that for some densities with high rate of decay and small variance like Exp(10) or Maxwell(20), the integrand $f(x) \log f(x)$ produces non-finite value for large value of x in R software due to the round-up. However, since the contribution of the right tail of the integrand to the entropy is insignificant for sufficiently large x , we obtain the approximately true value of entropy by cutting of the negligible right tail of the integration. That is, instead of integrating over the entire support of $f(x) \log f(x)$, we integrate up to a certain value at which the right tail is negligible.

3.1 Comparison of entropy estimators

In this sub-section, we run simulation experiments on different estimators which are classified into two groups, namely, ‘spacing estimators’ and ‘non-spacing estimators’.

3.1.1 Spacing estimators

1. Vasicek's estimator by Vasicek (1976):

$$\hat{H}_1 = \frac{1}{n} \sum_{i=1}^n \log \left\{ \frac{n}{2m} (X_{(i+m)} - X_{(i-m)}) \right\}. \quad (17)$$

Vasicek showed that if the variance is finite, then as $m \rightarrow \infty$ and $m/n \rightarrow 0$, $\hat{H}^{Vasicek}(X) \xrightarrow{\mathbb{P}} H(X)$.

2. Ebrahimi's estimator by Ebrahimi et al. (1992):

$$\hat{H}_2 = \frac{1}{n} \sum_{i=1}^n \log \left\{ \frac{n}{c_i m} (X_{(i+m)} - X_{(i-m)}) \right\}, \quad (18)$$

where

$$c_i = \begin{cases} 1 + \frac{i-1}{m} & 1 \leq i \leq m \\ 2 & m+1 \leq i \leq n-m \\ 1 + \frac{n-i}{m} & n-m+1 \leq i \leq n \end{cases}.$$

It was shown that $\hat{H}^{Ebrahimi}(X) \xrightarrow{\mathbb{P}} H(X)$ as $n, m \rightarrow \infty$, and $m/n \rightarrow 0$. Also, its bias is smaller than that of Vasicek's estimator.

3. Van's estimator by Van Es (1992):

$$\hat{H}_3 = -\frac{1}{n-m} \sum_{i=1}^{n-m} \log \left\{ \frac{n+1}{m} (X_{(i+m)} - X_{(i)}) \right\} + \sum_{k=m}^n \frac{1}{k} + \log \left(\frac{m}{n+1} \right). \quad (19)$$

Van established, under certain conditions, the strong consistency and asymptotic normality of $\hat{H}^{Van}(X)$.

4. Correa's estimator by Correa (1995):

$$\hat{H}_4 = -\frac{1}{n} \sum_{i=1}^{n-m} \log \left\{ \frac{\sum_{j=i-m}^{i+m} (X_{(j)} - \bar{X}_{(i)})(j-i)}{n \sum_{j=i-m}^{i+m} (X_{(j)} - \bar{X}_{(i)})} \right\}, \quad (20)$$

where $\bar{X}_{(i)} := \frac{1}{2m+1} \sum_{j=i-m}^{i+m} X_{(j)}$. The resulting estimator of entropy is shown to attain a smaller mean squared error (MSE) compared to that of Vasicek's estimator.

5. WG's estimator by by Wiczorkowski and Grzegorzewski (1999):

$$\hat{H}_5 = \hat{H}_1 + \log \left(\frac{2m}{n} \right) - \frac{n-2m}{n} \Psi(2m) + \Psi(n+1) - \frac{2}{n} \sum_{i=1}^m \Psi(i+m-1). \quad (21)$$

where $\Psi(k) = \sum_{i=1}^{k-1} \frac{1}{i} - \gamma$ is the di-Gamma function defined on integer set, and $\gamma = 0.57721566\dots$ is the Euler's constant.

6. Noughabi's estimator (kernel-spacing) by Noughabi (2010):

$$\hat{H}_6 = -\frac{1}{n} \sum_{i=1}^n \log \left\{ \frac{\hat{f}^{Fixed}(X_{(i+m)}) + \hat{f}^{Fixed}(X_{(i-m)})}{2} \right\}. \quad (22)$$

where the bandwidth in $\hat{f}^{Fixed}(x)$ is fixed to $b = 1.06sn^{-1/5}$ and s is the sample standard deviation. Similar to $\hat{H}^{Vasicek}(X)$, Noughabi showed that $\hat{H}^{Noughabi}(X)$ weakly converges to $H(X)$ using the same proof.

3.1.2 Non-spacing estimators

To see how the Poisson smoothed histogram density estimator performs in the estimation of entropy, we also run simulation on other direct plugin and mean of log entropy estimators that use the fixed symmetric kernel density estimator \hat{f}^{Fixed} and the Gamma asymmetric kernel density estimator \hat{f}^{Gam} which has the form

$$\hat{f}^{Gam}(x) = \frac{1}{n} \sum_{i=1}^n K_{x/b+1,b}^{Gam}(X_i) = \sum_{i=1}^n \frac{X_i^{x/b} \exp(-X_i/b)}{b^{x/b+1} \Gamma(x/b+1)}. \quad (23)$$

Thus, we obtain another four entropy estimators to compare with our proposed estimators. They are

$$\text{Plugin Fixed} : \hat{H}_7 = - \int_0^\infty \hat{f}^{Fixed}(x) \log \hat{f}^{Fixed}(x) dx, \quad (24)$$

$$\text{Meanlog Fixed} : \hat{H}_8 = -\frac{1}{n} \sum_{i=1}^n \log \hat{f}^{Fixed}(X_i), \quad (25)$$

$$\text{Plugin Gam} : \hat{H}_9 = - \int_0^\infty \hat{f}^{Gam}(x) \log \hat{f}^{Gam}(x) dx, \quad (26)$$

$$\text{Meanlog Gam} : \hat{H}_{10} = -\frac{1}{n} \sum_{i=1}^n \log \hat{f}^{Gam}(X_i). \quad (27)$$

The final entropy estimator we want to compare with is based on the quantile density estimator. Here the quantile function is first estimated by the method of Bernstein polynomial of degree m , which is of the form

$$\tilde{Q}_n(p) = \sum_{i=0}^m \hat{Q}_n\left(\frac{i}{m}\right) b(i, m, p) \quad p \in [0, 1],$$

where $\hat{Q}_n(\cdot)$ is the empirical quantile function, $b(i, m, p) = \mathbb{P}[Y = i]$ where Y follows the binomial(m, p), and m is a function of n such that $m \rightarrow \infty$ as $n \rightarrow \infty$. Then the estimator of $q(\cdot)$ can be obtained by differentiating $\tilde{Q}_n(\cdot)$

$$\tilde{q}_n(p) = \frac{d\tilde{Q}_n(p)}{dp} = \sum_{i=0}^m \hat{Q}_n\left(\frac{i}{m}\right) b(i, m, p) \left[\frac{i - mp}{p(1-p)} \right]. \quad (28)$$

Indeed, $\tilde{q}_n(\cdot)$ is the special case of the quantile kernel density estimator, proposed by Cheng and Parzen (1997), which has the general form

$$\tilde{q}_n^{CP}(p) := \frac{d}{dp} \tilde{Q}_n^{CP}(p) = \frac{d}{dp} \int_0^1 \hat{Q}_n(t) K_n(p, t) d\mu_n(t).$$

As a result, the last entropy estimator is of the form: $\hat{H}_{11} = \int_0^1 \log \tilde{q}_n(p) dp$. Together, we label our estimators as $\hat{H}_{12} = \hat{H}^{Plugin-Pois}$ and $\hat{H}_{13} = \hat{H}^{Meanlog-Pois}$.

The simulation study is organized as follows. For each density, 500 replicated data are generated for three sample sizes: $n = 10$, $n = 50$, and $n = 100$. To obtain the entropy estimator, we need to assign value to smoothing parameters. Particularly, since the optimal choice for m in spacing estimators is still an opening problem, for each spacing estimator, we compute the sample entropies for all values of m from 2 to $n/2 - 1$, then we use the optimal entropy with the smallest MSE to compare with other entropy estimators. On the other hand, the bandwidth selection in $\hat{f}^{Fixed}(\cdot)$ is set to $b = 1.06sn^{-1/5}$ while $b = sn^{-2/5}$ is used in $\hat{f}^{Gam}(\cdot)$. The choice of the polynomial degree m in the quantile density estimator $\tilde{q}(\cdot)$ is fixed to $m = n/\log n$. Lastly, for our estimators, we applied the smoothing parameter $k = n^{2/5} + 1$. To compare the performance between estimators, for each density and each estimator, we compute the point estimate and its MSE shown in the parentheses. The simulation results are shown in the Table ?? to Table ??. Each column in the table corresponds to one density with the associated true entropy right below, and the bold value in that column indicates the best estimator with the smallest MSE for that density. By observing the simulation results below, we have some important remarks.

Remark 3.1. *There does not exist the uniquely best entropy estimator in all cases.*

It is clear from the simulation results that depending on the density and the sample size, the best entropy estimator switches from one to another. When comparing entropy estimators, not only the smallest MSE is considered, but the computation expense is also an important key to determine the best estimator. Indeed, in terms of speed, the entropy estimators in the spacing group seem to dominate estimators in the other group. For instance for the sample size $n = 50$, it takes at most two seconds to obtain the spacing entropy estimators while it is up to thirty seconds for the direct plugin entropy estimator to be computed. This totally makes sense because the amount of computations in the latter estimators is much heavier than those in former estimators due to the presence of integration in their form. Definitely, this could be the main reason why most of papers in literature only focus on spacing estimators but a few in the direct plugin estimators.

		Monotone densities				
		Exp(1) H=1	Exp(10) H=-1.3026	Pareto(2,1) H=0.8069	Lnorm(0,2) H=2.1121	Weibull(0.5,0.5) H=0.4228
Spacing estimators	\hat{H}_1	0.5694(0.3167)	-1.7332(0.3167)	0.4857(0.3907)	1.8084(0.6484)	0.4196(0.4067)
	\hat{H}_2	0.8694(0.1505)	-1.4332(0.1505)	0.6350(0.2904)	2.0045(0.5677)	0.4785(0.3998)
	\hat{H}_3	0.8869(0.1503)	-1.4157(0.1503)	0.7083(0.2581)	1.9921(0.5311)	0.5426(0.4297)
	\hat{H}_4	0.7625(0.1896)	-1.5401(0.1896)	0.6311(0.2988)	2.0206(0.5815)	0.4936(0.4015)
	\hat{H}_5	0.9908(0.1313)	-1.3117(0.1313)	0.8515(0.2628)	2.2210(0.5680)	0.6950(0.4707)
	\hat{H}_6	1.0627(0.1428)	-1.2399(0.1428)	1.1180(0.5314)	2.9613(1.8878)	1.2738(1.2629)
Non-spacing estimators	\hat{H}_7	1.0142(0.0704)	divergent	0.7446(0.0659)	1.2540(0.9033)	divergent
	\hat{H}_8	1.0934(0.1525)	-1.2091(0.1525)	1.1658(0.5780)	3.0210(2.0037)	1.3304(1.3736)
	\hat{H}_9	1.0390(0.0842)	divergent	1.1095(0.2202)	1.7242(0.3123)	0.9027(0.3854)
	\hat{H}_{10}	1.0102(0.1178)	-1.2923(0.1178)	1.2326(0.4528)	2.5934(1.0828)	0.9867(0.7638)
	\hat{H}_{11}	0.7272(0.2140)	1.5754(0.2140)	0.6924(0.3503)	2.2866(0.8415)	0.6751(0.5048)
	\hat{H}_{12}	1.0610(0.0676)	-0.2004(1.2198)	1.1612(0.1958)	1.7180(0.3854)	0.8643(0.3612)
	\hat{H}_{13}	0.8677(0.0983)	-0.8690(0.2064)	1.0988(0.1748)	1.6043(0.5424)	0.4957(0.2262)

Table 1: Simulation results for $n = 10$ of the monotone density group. The point estimate and MSE (in parentheses) are computed for each estimator. The bold value is the best estimator with the smallest MSE.

		Unimodal densities					
		Gam(2,2) H=0.8841	Gam(7.5,1) H=2.3804	Lnorm(0,0.5) H=0.7258	Maxwell(1) H=0.9962	Maxwell(20) H=0.5017	Weibull(2,2) H=1.2886
Spacing estimators	\hat{H}_1	0.3676(0.3560)	1.8444(0.3569)	0.2076(0.2723)	0.4790(0.3319)	-1.0189(0.3319)	0.7885(0.3228)
	\hat{H}_2	0.6342(0.1489)	2.0746(0.1576)	0.4608(0.1746)	0.7084(0.1407)	-0.7895(0.1407)	1.0229(0.1371)
	\hat{H}_3	0.7103(0.1300)	2.1784(0.1191)	0.5316(0.1482)	0.8290(0.1045)	-0.6688(0.1045)	1.1352(0.1096)
	\hat{H}_4	0.5401(0.2063)	2.0139(0.2043)	0.3786(0.2259)	0.6481(0.1860)	-0.8498(0.1860)	0.9577(0.1822)
	\hat{H}_5	0.7745(0.0973)	2.2571(0.0848)	0.6203(0.1149)	0.8917(0.0754)	-0.6062(0.0754)	1.1787(0.0801)
	\hat{H}_6	0.9023(0.0874)	2.3403(0.0646)	0.7244(0.1042)	0.9756(0.0588)	-0.5223(0.0588)	1.2893(0.0668)
Non-spacing estimators	\hat{H}_7	0.8891(0.0613)	2.4273(0.0548)	0.8021(0.0895)	1.0301(0.0435)	divergent	1.2912(0.0442)
	\hat{H}_8	0.8332(0.0951)	2.2495(0.0802)	0.6503(0.1156)	0.8840(0.0711)	-0.6139(0.0711)	1.2016(0.0756)
	\hat{H}_9	0.9395(0.0571)	2.6388(0.0968)	0.9217(0.0959)	1.1949(0.0648)	-0.2888(0.0731)	1.4138(0.0467)
	\hat{H}_{10}	0.8406(0.0742)	2.3963(0.0444)	0.7333(0.0801)	0.9933(0.0395)	-0.5046(0.0395)	1.2688(0.0489)
	\hat{H}_{11}	0.5040(0.2332)	1.9697(0.2387)	0.3450(0.2533)	0.5998(0.2212)	0.8980(0.2212)	0.9114(0.2112)
	\hat{H}_{12}	1.0245(0.0532)	2.3802(0.0257)	1.0476(0.1271)	1.2644(0.0832)	0.2563(0.5828)	1.4288(0.0387)
	\hat{H}_{13}	0.8120(0.0504)	2.1908(0.0707)	0.7665(0.0400)	1.0046(0.0215)	-0.1038(0.1655)	1.2183(0.0353)

Table 2: Simulation results for $n = 10$ of the unimodal density group. The point estimate and MSE (in parentheses) are computed for each estimator. The bold value is the best estimator with the smallest MSE.

		Bimodal densities			
		MixGam H=2.2757	MixLnorm H=1.6724	MixMaxwell H=0.8014	MixWeibull H=1.1330
Spacing estimators	\hat{H}_1	2.0233(0.0945)	1.3541(0.5505)	0.4218(0.1888)	0.9726(0.1635)
	\hat{H}_2	2.3402(0.0350)	1.5502(0.4641)	0.7387(0.0486)	1.2228(0.1480)
	\hat{H}_3	2.4074(0.0504)	1.4707(0.3565)	0.8033(0.0595)	1.3422(0.1414)
	\hat{H}_4	2.2469(0.0333)	1.5604(0.4861)	0.6319(0.0734)	1.1713(0.1532)
	\hat{H}_5	2.3214(0.0432)	1.7667(0.4581)	0.8247(0.0444)	1.3253(0.1918)
	\hat{H}_6	2.6307(0.1614)	2.2060(1.5798)	0.9993(0.0913)	1.5081(0.2682)
Non-spacing estimators	\hat{H}_7	2.4427(0.0539)	1.2077(0.3345)	1.0057(0.0703)	1.3375(0.0623)
	\hat{H}_8	2.4700(0.0946)	2.2567(1.6634)	1.0161(0.0967)	1.5286(0.1927)
	\hat{H}_9	2.5721(0.1097)	1.4677(0.1810)	1.0041(0.0675)	1.3452(0.0695)
	\hat{H}_{10}	2.5106(0.0952)	1.9468(0.8679)	0.9495(0.0585)	1.3924(0.1293)
	\hat{H}_{11}	2.1290(0.0563)	1.7298(0.8283)	0.5777(0.0961)	1.0935(0.1563)
	\hat{H}_{12}	2.2509(0.0190)	1.5151(0.1883)	1.0488(0.0793)	1.3312(0.0664)
	\hat{H}_{13}	2.0440(0.0813)	1.3316(0.2847)	0.8673(0.0264)	1.1107(0.1077)

Table 3: Simulation results for $n = 10$ of the bimodal density group. The point estimate and MSE (in parentheses) are computed for each estimator. The bold value is the best estimator with the smallest MSE.

		Monotone densities				
		Exp(1) H=1	Exp(10) H=-1.3026	Pareto(2,1) H=0.8069	Lnorm(0,2) H=2.1121	Weibull(0.5,0.5) H=0.4228
Spacing estimators	\hat{H}_1	0.8818(0.0368)	-1.4207(0.0368)	0.7405(0.0663)	2.0568(0.1184)	0.4555(0.1011)
	\hat{H}_2	0.9941(0.0229)	-1.3085(0.0229)	0.7653(0.0587)	2.0646(0.1143)	0.4475(0.1029)
	\hat{H}_3	0.9516(0.0270)	-1.3509(0.0270)	0.7454(0.0547)	2.0646(0.1123)	0.4677(0.1002)
	\hat{H}_4	0.9758(0.0236)	-1.3268(0.0236)	0.7875(0.0570)	2.0793(0.1120)	0.4595(0.1054)
	\hat{H}_5	1.0174(0.0230)	-1.2851(0.0230)	0.8142(0.0535)	2.1403(0.1111)	0.4647(0.1065)
	\hat{H}_6	1.1124(0.0364)	-1.1902(0.0364)	1.2279(0.3808)	3.4125(2.2625)	1.3432(0.9828)
Non-spacing estimators	\hat{H}_7	1.3702(0.1554)	divergent	1.0947(0.1223)	2.1172(0.0657)	1.2002(0.6431)
	\hat{H}_8	1.5531(0.3478)	-1.1404(0.0506)	1.3116(0.4532)	3.5042(2.4923)	1.4557(1.1993)
	\hat{H}_9	1.0648(0.0219)	-1.0389(0.0915)	1.1494(0.1534)	1.7962(0.1410)	1.0498(0.4362)
	\hat{H}_{10}	1.0577(0.0250)	-1.2449(0.0250)	1.2538(0.2882)	2.8416(0.8425)	1.0126(0.4482)
	\hat{H}_{11}	1.0215(0.0250)	1.2811(0.0250)	0.9867(0.1284)	2.5532(0.3923)	0.7510(0.2102)
	\hat{H}_{12}	1.0665(0.0215)	-0.5605(0.5543)	1.1189(0.1188)	1.8701(0.1160)	0.7985(0.1898)
	\hat{H}_{13}	0.9650(0.0211)	-1.0823(0.0581)	1.0811(0.1031)	1.9167(0.1194)	0.5376(0.0785)

Table 4: Simulation results for $n = 50$ of the monotone density group. The point estimate and MSE (in parentheses) are computed for each estimator. The bold value is the best estimator with the smallest MSE.

		Unimodal densities					
		Gam(2,2) H=0.8841	Gam(7.5,1) H=2.3804	Lnorm(0,0.5) H=0.7258	Maxwell(1) H=0.9962	Maxwell(20) H=-0.5017	Weibull(2,2) H=1.2886
Spacing estimators	\hat{H}_1	0.7377(0.0363)	2.2204(0.0379)	0.5805(0.0419)	0.8436(0.0320)	-0.6543(0.0320)	1.1291(0.0361)
	\hat{H}_2	0.8597(0.0165)	2.3436(0.0151)	0.6915(0.0238)	0.9655(0.0103)	-0.5324(0.0103)	1.2480(0.0135)
	\hat{H}_3	0.8264(0.0214)	2.3025(0.0216)	0.6508(0.0279)	0.9360(0.0157)	-0.5619(0.0157)	1.2244(0.0172)
	\hat{H}_4	0.8327(0.0178)	2.3166(0.0166)	0.6756(0.0234)	0.9398(0.0123)	-0.5581(0.0123)	1.2238(0.0153)
	\hat{H}_5	0.8792(0.0149)	2.3619(0.0126)	0.7183(0.0206)	0.9869(0.0087)	-0.5109(0.0087)	1.2707(0.0109)
	\hat{H}_6	0.8962(0.0164)	2.3811(0.0124)	0.7300(0.0194)	0.9848(0.0102)	-0.5131(0.0102)	1.2865(0.0110)
Non-spacing estimators	\hat{H}_7	0.9466(0.0164)	2.469(0.0188)	0.5744(0.0347)	1.0689(0.0126)	divergent	1.3212(0.0088)
	\hat{H}_8	0.9191(0.0175)	2.3518(0.0125)	0.7390(0.0221)	0.9785(0.0093)	-0.5194(0.0093)	1.2748(0.0107)
	\hat{H}_9	0.9593(0.0171)	2.5833(0.0489)	0.9091(0.0492)	1.1594(0.0323)	-0.3360(0.0332)	1.3908(0.0172)
	\hat{H}_{10}	0.8970(0.0139)	2.4078(0.0105)	0.7630(0.0195)	1.0153(0.0079)	-0.4825(0.0079)	1.2899(0.0089)
	\hat{H}_{11}	0.8705(0.0169)	2.3480(0.0151)	0.7267(0.0259)	0.9621(0.0105)	-0.5358(0.0105)	1.2434(0.0138)
	\hat{H}_{12}	1.0109(0.0247)	2.4258(0.0096)	0.9796(0.0726)	1.2045(0.0468)	0.1018(0.3659)	1.3998(0.0176)
	\hat{H}_{13}	0.8797(0.0113)	2.3276(0.0122)	0.7750(0.0150)	1.0245(0.0067)	-0.2296(0.0765)	1.2728(0.0078)

Table 5: Simulation results for $n = 50$ of the unimodal density group. The point estimate and MSE (in parentheses) are computed for each estimator. The bold value is the best estimator with the smallest MSE.

		Bimodal densities			
		MixGam H=2.2757	MixLnorm H=1.6724	MixMaxwell H=0.8014	MixWeibull H=1.1330
Spacing estimators	\hat{H}_1	2.2335(0.0117)	1.6370(0.0742)	0.6712(0.0237)	1.1196(0.0440)
	\hat{H}_2	2.2732(0.0086)	1.6386(0.0707)	0.7997(0.0069)	1.1524(0.0431)
	\hat{H}_3	2.2847(0.0080)	1.5939(0.0704)	0.7801(0.0100)	1.1969(0.0306)
	\hat{H}_4	2.2792(0.0088)	1.6486(0.0684)	0.7649(0.0082)	1.1160(0.0438)
	\hat{H}_5	2.2955(0.0099)	1.7084(0.0690)	0.8109(0.0067)	1.1781(0.0444)
	\hat{H}_6	2.5927(0.1088)	2.6941(1.7221)	0.9969(0.0462)	1.5212(0.1882)
Non-spacing estimators	\hat{H}_7	2.5123(0.0619)	1.5856(0.0733)	1.0167(0.0515)	1.3733(0.0263)
	\hat{H}_8	2.5520(0.0864)	2.7858(1.9013)	1.0072(0.0503)	1.5531(0.1118)
	\hat{H}_9	2.6064(0.1150)	1.6203(0.0302)	0.9806(0.0369)	1.3559(0.0239)
	\hat{H}_{10}	2.5520(0.0845)	2.1751(0.5432)	0.9149(0.0185)	1.3840(0.0428)
	\hat{H}_{11}	2.3299(0.0121)	2.1617(0.4259)	0.8168(0.0079)	1.3226(0.0996)
	\hat{H}_{12}	2.3168(0.0077)	1.5664(0.0495)	1.0213(0.0522)	1.3105(0.0195)
	\hat{H}_{13}	2.2166(0.0118)	1.5316(0.0676)	0.8761(0.0097)	1.1669(0.0378)

Table 6: Simulation results for $n = 50$ of the bimodal density group. The point estimate and MSE (in parentheses) are computed for each estimator. The bold value is the best estimator with the smallest MSE.

		Monotone densities				
		Exp(1) H=1	Exp(10) H=-1.3026	Pareto(2,1) H=0.8069	Lnorm(0,2) H=2.1121	Weibull(0.5,0.5) H=0.4228
Spacing estimators	\hat{H}_1	0.9246(0.0166)	-1.3780(0.0166)	0.7705(0.0290)	2.0973(0.0518)	0.4337(0.0466)
	\hat{H}_2	0.9975(0.0107)	-1.3051(0.0107)	0.7820(0.0265)	2.1137(0.0509)	0.4323(0.0469)
	\hat{H}_3	0.9595(0.0129)	-1.3431(0.0129)	0.7715(0.0256)	2.0754(0.0521)	0.4448(0.0455)
	\hat{H}_4	0.9854(0.0107)	-1.3172(0.0107)	0.8010(0.0250)	2.1238(0.0507)	0.4409(0.0479)
	\hat{H}_5	1.0030(0.0104)	-1.2996(0.0104)	0.8152(0.0244)	2.1296(0.0504)	0.4469(0.0485)
	\hat{H}_6	1.0916(0.0196)	-1.2110(0.0196)	1.2405(0.3582)	3.4532(2.1463)	1.3183(0.8771)
Non-spacing estimators	\hat{H}_7	1.0980(0.0175)	divergent	1.1411(0.1391)	2.1739(0.0389)	1.2275(0.6685)
	\hat{H}_8	1.1501(0.0335)	-1.1525(0.0335)	1.3429(0.4410)	3.5585(2.4107)	1.4594(1.1432)
	\hat{H}_9	1.0501(0.0111)	-1.0950(0.0534)	1.1478(0.1347)	1.8295(0.1033)	1.0531(0.4198)
	\hat{H}_{10}	1.0451(0.0121)	-1.2575(0.0121)	1.2469(0.2491)	2.8244(0.6626)	0.9872(0.3663)
	\hat{H}_{11}	1.0375(0.0128)	-1.2651(0.0128)	0.6637(0.0502)	1.9686(0.0903)	divergent
	\hat{H}_{12}	1.0041(0.0066)	-0.6973(0.3688)	1.2950(0.2498)	1.8994(0.0776)	0.7686(0.1453)
	\hat{H}_{13}	0.9733(0.0103)	-1.1530(0.0280)	1.0713(0.0851)	1.9845(0.0585)	0.5433(0.0476)

Table 7: Simulation results for $n = 100$ of the monotone density group. The point estimate and MSE (in parentheses) are computed for each estimator. The bold value is the best estimator with the smallest MSE.

		Unimodal densities					
		Gam(2,2) H=0.8841	Gam(7.5,1) H=2.3804	Lnorm(0,0.5) H=0.7258	Maxwell(1) H=0.9962	Maxwell(20) H=-0.5017	Weibull(2,2) H=1.2886
Spacing estimators	\hat{H}_1	0.7951(0.0150)	2.2896(0.0147)	0.6469(0.0151)	0.9033(0.0132)	-0.5946(0.0132)	1.193(0.0140)
	\hat{H}_2	0.8742(0.0075)	2.3668(0.0060)	0.7153(0.0090)	0.9882(0.0049)	-0.5096(0.0049)	1.2763(0.0055)
	\hat{H}_3	0.8446(0.0102)	2.3333(0.0097)	0.6770(0.0125)	0.9569(0.0077)	-0.5410(0.0077)	1.2508(0.0080)
	\hat{H}_4	0.8629(0.0076)	2.3587(0.0071)	0.7106(0.0088)	0.9732(0.0053)	-0.5247(0.0053)	1.2635(0.0056)
	\hat{H}_5	0.8843(0.0070)	2.3777(0.0064)	0.7280(0.0085)	0.9983(0.0046)	-0.4995(0.0046)	1.2851(0.0049)
	\hat{H}_6	0.8939(0.0079)	2.3560(0.0068)	0.7296(0.0087)	0.9779(0.0049)	-0.5200(0.0049)	1.2794(0.0048)
Non-spacing estimators	\hat{H}_7	0.9397(0.0093)	2.4614(0.0124)	0.8477(0.0235)	1.0597(0.0079)	divergent	1.3233(0.0049)
	\hat{H}_8	0.9180(0.0089)	2.3663(0.0063)	0.7400(0.0093)	0.9866(0.0046)	-0.5113(0.0046)	1.2873(0.0047)
	\hat{H}_9	0.9464(0.0097)	2.5499(0.0331)	0.8808(0.0310)	1.1301(0.0210)	-0.3666(0.0214)	1.3782(0.0114)
	\hat{H}_{10}	0.8930(0.0068)	2.4025(0.0059)	0.7537(0.0086)	1.0094(0.0041)	-0.4885(0.0041)	1.2935(0.0042)
	\hat{H}_{11}	0.8628(0.0100)	2.3522(0.0103)	0.6953(0.0137)	0.9700(0.0073)	0.5278(0.0073)	1.2546(0.0079)
	\hat{H}_{12}	0.9891(0.0156)	2.4262(0.0065)	0.9407(0.0503)	1.1704(0.0324)	0.0272(0.2806)	1.3856(0.0121)
	\hat{H}_{13}	0.8816(0.0058)	2.3514(0.0063)	0.7639(0.0076)	1.0180(0.0038)	-0.2849(0.0485)	1.2834(0.0038)

Table 8: Simulation results for $n = 100$ of the unimodal density group. The point estimate and MSE (in parentheses) are computed for each estimator. The bold value is the best estimator with the smallest MSE.

		Bimodal densities			
		MixGam H=2.2757	MixLnorm H=1.6724	MixMaxwell H=0.8014	MixWeibull H=1.1330
Spacing estimators	\hat{H}_1	2.2615(0.0054)	1.6434(0.0346)	0.7194(0.0103)	1.1248(0.0216)
	\hat{H}_2	2.2818(0.0037)	1.6476(0.0335)	0.7998(0.0037)	1.1424(0.0213)
	\hat{H}_3	2.2834(0.0033)	1.6211(0.0348)	0.7926(0.0053)	1.1735(0.0154)
	\hat{H}_4	2.2817(0.0037)	1.6850(0.0327)	0.7854(0.0040)	1.1513(0.0215)
	\hat{H}_5	2.2859(0.0041)	1.6869(0.0326)	0.8035(0.0036)	1.1524(0.0214)
	\hat{H}_6	2.5557(0.0824)	2.8079(1.8217)	0.9702(0.0328)	1.5147(0.1646)
Non-spacing estimators	\hat{H}_7	2.5170(0.0611)	1.8478(0.0672)	1.0020(0.0432)	1.3790(0.0175)
	\hat{H}_8	2.5496(0.0795)	2.9098(2.0306)	0.9793(0.0359)	1.5534(0.0919)
	\hat{H}_9	2.5966(0.1058)	1.6365(0.0158)	0.9584(0.0273)	1.3569(0.0144)
	\hat{H}_{10}	2.537(0.0755)	2.1910(0.4629)	0.8873(0.0104)	1.3706(0.0239)
	\hat{H}_{11}	2.4534(0.0393)	1.5709(0.0510)	0.7508(0.0076)	Divergent
	\hat{H}_{12}	2.3167(0.0044)	1.5792(0.0287)	0.9955(0.0398)	1.2996(0.0100)
	\hat{H}_{13}	2.2402(0.0050)	1.5759(0.0345)	0.8617(0.0059)	1.1753(0.0250)

Table 9: Simulation results for $n = 100$ of the bimodal density group. The point estimate and MSE (in parentheses) are computed for each estimator. The bold value is the best estimator with the smallest MSE.

Remark 3.2. *The direct plugin entropy estimators by means of \hat{f}^{Fixed} , \hat{f}^{Gam} , and $\hat{q}^{Quantile}$ encounter the convergence problem when computing the integral.*

Another issue when we run the direct plugin entropy estimators is the convergence of the integral. Perhaps, this is the greatest weakness of the direct plugin estimators. Indeed, in most cases, we could not evaluate the integral in $\hat{H}_7 = \hat{H}^{Plugin-Fixed}(X)$ and $\hat{H}_9 = \hat{H}^{Plugin-Gam}(X)$ over the same interval that we used to compute the true entropy because it produces non-finite values on the right tail. Thus, we have to cut off the larger tail of the integral to a shorter finite interval up to the point that the value of the integrand is finite. This issue, however, does not appear in our entropy estimator $\hat{H}_{12} = \hat{H}^{Plugin-Pois}$, so it makes our entropy estimator more favorable than other entropy estimators based on the direct plugin approach.

Remark 3.3. *Our entropy estimator by the mean of log of Poisson distribution based density estimator $\hat{H}_{13} = \hat{H}^{Meanlog-Pois}$ seems to be the best entropy estimator when the sample size is small.*

We see that most of the times, the MSE of \hat{H}_{13} is the smallest comparing to that of others estimators for small sample size in the same density. Note that we have to take into account that while the spacing

entropy estimators are computed with the best optimal spacing parameter m , the smoothing parameter in our estimators is set to $k = n^{2/5} + 1$, which may be far away from the optimal value. Still, our entropy estimator \hat{H}_{13} outperforms other entropy estimators in most cases.

Remark 3.4. *As the sample size increases, the spacing entropy estimators, \hat{H}_2 , \hat{H}_3 , \hat{H}_4 , and \hat{H}_5 , perform much better, and converge to the true value.*

Although for a small sample size like $n = 10$ (Table ??-??), the performance of the entropy estimators from spacing group is quite poor comparing to the estimators in other group, their rate of convergence, however, pick up very fast as the sample size increases as in $n = 100$. For instance, the number of the best entropy estimator that falls into the spacing group increase as the sample size increases from 10 to 100. Therefore, together with the fast computation, the spacing entropy estimators would be a better choice in the case of large sample size.

In conclusion, there is a trade-off between the speed and the performance for the two groups of entropy estimators. It is obvious that any spacing estimator (Ebrihami's estimator, Van Es estimator, Correa's estimator, or WG estimator) would do a good job when dealing with large sample sizes due to its fast rate of convergence and fast computation; though the answer to the question of what value of m would be considered for a large sample size is still unknown. However, when sample size is small, the choice of estimator for entropy becomes more difficult and requires careful studies. This is why in the small sample size, the entropy estimators by means of density estimator come to handy because they give a more precise result. Among non-spacing entropy estimators presented above, our mean of log of Poisson density points \hat{H}_{13} consistently performs better than the others.

3.2 Residual entropy estimators comparison

In this subsection, we want to see the performance of our proposed estimator for residual entropy function as well as comparing it with the existing ones. Particularly, we still use the same set of testing distributions in the previous section, and the set of estimators are

$$\begin{aligned} \hat{H}_1^{Belzunce}(X, t) &= \log \hat{R}(t) - \frac{1}{\hat{R}(t)} \int_t^\infty \hat{f}^{Fixed}(x) \log \hat{f}^{Fixed}(x) dx, \\ \hat{H}_2^{Belzunce}(X, t) &= \log \hat{R}(t) - \frac{1}{\hat{R}(t)} \sum_{i=1}^n R_K \left(\frac{t - X_i}{b} \right) \log \hat{f}_i^{Fixed}(X_i), \\ \hat{H}^{Quantile}(X, t) &= \log(\hat{R}^{Pois}(t)) - \frac{1}{\hat{R}^{Pois}(t)} \int_{\hat{F}(t)}^1 \log \tilde{q}_n(p) dp, \\ \hat{H}^{Pois}(X, t) &= \log \hat{R}^{Pois}(t) - \frac{1}{\hat{R}^{Pois}(t)} \int_t^\infty \hat{f}^{Pois}(x) \log \hat{f}^{Pois}(x) dx. \end{aligned}$$

where the estimated survival function $\hat{R}(t)$ is computed based on the integration of $\hat{f}^{Fixed}(\cdot)$, the bandwidth is fixed to $b = 1.06sn^{-1/5}$ in \hat{f}^{Fixed} , the smoothing parameter k in \hat{f}^{Pois} is assigned to $n^{2/5} + 1$, and the parameter m in $\tilde{q}_n(\cdot)$ is set equal to $n/\log n$.

The setup of simulation study is as following. We present here both performance comparison by graphics (for $n = 50$) and by mean intergrated squared error (MISE) (for $n = 50, 100$ and 500). In order to produce the plots of residual entropy estimators, for each distribution, we run 500 replications, then

we compute the sample mean residual entropy estimator functions of these 500 replications. As mentioned in Belzunce et al. (2001) that the behavior of residual entropy function is smooth up to certain bound because $R(t) \rightarrow 0$ very fast when $t \rightarrow \infty$, it is recommended to stop the estimation at the time $t^* = \inf\{t : R(t) \leq 0.01\}$. Therefore, the MISEs are just computed on the interval $[0, t^*]$ (for Pareto(2,1) is $[1, t^*]$). Similarly, the MISEs comparison between estimators are done based on 500 replications for sample size $n = 50$ and $n = 100$ for each distribution, but only 100 replications for sample size $n = 500$. The plots and tables below show the performance of our estimators and the ones in Belzunce et al. (2001).




Figure 1: Plots of residual entropy estimators for $n = 50$. The true function is in black solid line, $\hat{H}^{Pois}(X, t)$ in red dotted-dashed line, $\hat{H}^{Quantile}(X, t)$ in black long dashed line, $\hat{H}^{Belzunce}(X, t)$ in blue dashed line, and $\hat{H}_2^{Belzunce}(X, t)$ in green dotted line. There are some plots without the green dotted line because the function is not in the range.




Figure 2: Plots of residual entropy estimators for $n = 50$. The true function is in black solid line, $\hat{H}^{Pois}(X, t)$ in red dotted-dashed line, $\hat{H}^{Quantile}(X, t)$ in black long dashed line, $\hat{H}_1^{Belzunce}(X, t)$ in blue dashed line, and $\hat{H}_2^{Belzunce}(X, t)$ in green dotted line.

Intergrated Mean Squared Error (IMSE)

	Interval	$\hat{H}^{Pois}(X, t)$	$\hat{H}_1^{Belzunce}(X, t)$	$\hat{H}^{Quantile}(X, t)$
Exp(1)	(0,4.605)	0.608	2.5816	9.8387
Exp(10)	(0,0.461)	0.2535	1.3814	0.0719
Gamma(7.5,1)	(0,15.289)	0.9315	1.5832	14.4769
Gamma(2,2)	(0,3.319)	0.1993	0.9969	3.1009
LogNormal(0,2)	(0,3.200)	2.2236	0.9033	2.9778
LogNormal(0,0.5)	(0,3.200)	0.2631	2.188	3.1081
Weibull(0.5,0.5)	(0,10.604)	15.6378	>10000	>10000
Weibull(2,2)	(0,4.292)	0.2945	0.3276	1.797
Maxwell(1)	(0,3.368)	0.4408	0.1853	0.6615
Maxwell(20)	(0,0.753)	0.7173	0.0379	0.0367
Pareto(2)	(1,10.024)	19.6957	6509.037	181.6876
MixGamma	(0,14.130)	1.1988	0.8582	17.3737
MixLogNormal	(0,3.200)	2.5622	3.2082	3.6732
MixWeibull	(0,4.292)	2.3112	0.7388	3.9595
MixMaxwell	(0,3.136)	0.365	0.1949	0.9866

Table 10: MISEs of three estimators on 500 replications with sample size $n = 50$. The MISEs are computed on the indicated interval for each distribution.

Intergrated Mean Squared Error (IMSE)				
	Interval	$\hat{H}^{Pois}(X,t)$	$\hat{H}_1^{Belzunce}(X,t)$	$\hat{H}^{Quantile}(X,t)$
Exp(1)	(0,4.605)	0.4124	2.4588	4.5337
Exp(10)	(0,0.461)	0.1674	0.2396	0.0362
Gamma(7.5,1)	(0,15.289)	0.6	0.944	5.9302
Gamma(2,2)	(0,3.319)	0.1505	0.6284	1.5077
LogNormal(0,2)	(0,3.200)	0.7871	0.9182	1.7078
LogNormal(0,0.5)	(0,3.200)	0.1826	0.8065	1.4114
Weibull(0.5,0.5)	(0,10.604)	9.5602	40.3328	62.3692
Weibull(2,2)	(0,4.292)	0.2113	0.2012	0.7826
Maxwell(1)	(0,3.368)	0.3158	0.1346	0.3129
Maxwell(20)	(0,0.753)	0.5532	0.03	0.031
Pareto(2)	(1,10.024)	13.5616	26773.74	66.4542
MixGamma	(0,14.130)	0.7175	0.8547	6.9898
MixLogNormal	(0,3.200)	1.0251	2.8335	2.8198
MixWeibull	(0,4.292)	2.5913	0.9241	4.6132
MixMaxwell	(0,3.136)	0.258	0.2852	0.4339

Table 11: MISEs of three estimators on 500 replications with sample size $n = 100$. The MISEs are computed on the indicated interval for each distribution.

Intergrated Mean Squared Error (IMSE)				
	Interval	$\hat{H}^{Pois}(X,t)$	$\hat{H}_1^{Belzunce}(X,t)$	$\hat{H}^{Quantile}(X,t)$
Exp(1)	(0,4.605)	0.1013	0.7366	0.4834
Exp(10)	(0,0.461)	0.0583	0.06	0.0099
Gamma(7.5,1)	(0,15.289)	0.1677	0.3434	0.5875
Gamma(2,2)	(0,3.319)	0.0537	0.1211	0.1674
LogNormal(0,2)	(0,3.200)	0.3641	0.7894	0.2377
LogNormal(0,0.5)	(0,3.200)	0.0722	0.1486	0.2194
Weibull(0.5,0.5)	(0,10.604)	1.8837	28.6434	3.6078
Weibull(2,2)	(0,4.292)	0.0843	0.0597	0.124
Maxwell(1)	(0,3.368)	0.1273	0.0356	0.0548
Maxwell(20)	(0,0.753)	0.2719	0.0066	0.0181
Pareto(2)	(1,10.024)	3.3839	>1000	3.4142
MixGamma	(0,14.130)	0.1736	0.8307	0.5792
MixLogNormal	(0,3.200)	0.4343	1.9141	0.5238
MixWeibull	(0,4.292)	3.02	0.8817	6.2564
MixMaxwell	(0,3.136)	0.103	0.1164	0.0786

Table 12: MISEs of three estimators on 100 replications with sample size $n = 500$. The MISEs are computed on the indicated interval for each distribution.

To illustrate the rate of convergence of each estimators as the sample size increases, we also show the following graphs which are MISEs as a function of sample size for each distribution.




Figure 3: Residual entropy estimation MISE comparison with different sample size $n = 50, 100$ and 500 . Where diamond-red is $\hat{H}^{Pois}(X,t)$, square-blue is $\hat{H}^{1Belzunce}(X,t)$, and triangle-black is $\hat{H}^{Quantile}(X,t)$.




Figure 4: Residual entropy estimation MISE comparison with different sample size $n = 50, 100$ and 500 . Where diamond-red is $\hat{H}^{Pois}(X, t)$, square-blue is $\hat{H}_1^{Belzunce}(X, t)$, and triangle-black is $\hat{H}^{Quantile}(X, t)$.

Remark 3.5. $\hat{H}_2^{Belzunce}(X, t)$ performs very poor in most cases.

By the plot of residual entropy estimations from Figure ??-??, we see that $\hat{H}_2^{Belzunce}(X, t)$ is not as good as the others among all estimators. It is even out of the range of MISE in some distributions. Therefore, we decide to remove it out of our consideration.

Remark 3.6. All estimators tend to diverge away from the true residual entropy over the time in most cases.

We notice that most of estimators have a quite good startup as t is small. However, as the time t increases, in which the true survival function tends to zero rapidly, the estimators begin to diverge away from the true residual entropy. How large of the time t this phenomenon happens depends on the true distribution.

Remark 3.7. None of estimators outperforms the others in all cases. However, the $\hat{H}^{Pois}(X, t)$ estimator seems to achieve a better estimation in most cases.

If we just focus on the performance comparison between estimators, we see that in most cases, our estimator $\hat{H}^{Pois}(X, t)$ captures the trend of the true function throughout the plots and achieves better precision in terms of MISE shown in the Table ??-??. It is defeated by $\hat{H}_1^{Belzunce}(X, t)$ only in Maxwell distributions which have the bell-normal shape. Therefore, the $\hat{H}^{Pois}(X, t)$ is recommended in a small sample size.

Remark 3.8. The rate of convergence of the estimator $\hat{H}^{Quantile}(X, t)$ seems to be faster than other competitors.

Similar to the entropy estimation, the residual entropy estimator, $\hat{H}^{Quantile}(X, t)$ still has the fastest rate of convergence in most cases as the sample size increases. We admit that for a small sample size, $\hat{H}^{Quantile}(X, t)$ does not show up to be a good estimator, and it is not recommended. However, if the sample size is sufficiently large, it becomes one of the best potential choices for residual entropy estimator.

References

- [1] Belzunce, F., Guillamon, A., Navarro, J., and Ruiz, J. M. (2001) Kernel estimation of residual entropy. *Communications in Statistics-Theory and Methods* 30:1243-1255.
- [2] Bouzebda, S., Elhattab, I., Keziou, A., and Lounis, T. (2013) New entropy estimator with an application to test of normality. *Communication in Statistics-Theory and Methods* 42:2245-2270.
- [3] Bouezmarni, T. and Scaillet, O. (2005) Consistency of asymmetric kernel density estimators and smoothed histograms with application to income data. *Econometric Theory* 21:390-412.
- [4] Chaubey, Y. P., Mudholkar, G. S. and Smethurst, P. A. (1993). On entropy-based goodness-of-fit tests: a practical strategy. In *Probability and Statistics*, S. K. Basu and B. K. Sinha, eds., Narosa Publishing House, New Delhi, 116-120.
- [5] Chaubey, Y. P. and Sen, P. K. (2009). On The Selection Of The Smoothing Parameter In Poisson Smoothing Of Histogram Estimator: Computational Aspects. *Pakistan Journal of Statistics* **25(4)**: 385-401.
- [6] Chaubey, Y. P., Sen, P., and Li, J. (2010) Smooth density estimation for length-biased data. *Journal of The Indian Society of Agricultural Statistics* 64(2):145-155.
- [7] Cheng, C. and Parzen, E. (1997) Unified estimators of smooth quantile and quantile density functions. *Journal of Statistical Planning and Inference* 59:291-307.
- [8] Correa, J. C. (1995). A new estimator of entropy. *Communications in Statistics: Theory and Methods* 24:2439-2449.
- [9] Cover, T. M. and Thomas J. A. (1991). *Elements of Information Theory*. Wiley, New York, USA.
- [10] Ebrahimi, N., Habibullah, M., and Soofi, E. (1992) *Testing exponentiality based on KullbackLeibler information*. Journal of the Royal Statistical Society: Series B, 54:739-748.
- [11] Ebrahimi, N. (1996). How to measure uncertainty in the life time distributions. *The Indian Journal of Statistics Series A* 58:48-56.
- [12] Eggermont, P.B. and LaRiccia V. N. (1999). Best Asymptotic Normality of the Kernel Density Entropy Estimator for Smooth Densities. *IEEE Transactions On Information Theory* 45:1321-1326
- [13] Hall, P. and Morton, S. C. (1993). On the estimation of entropy. *Ann. Inst. Statist. Math.* 45:69-88.
- [14] Kapur, J. N. (1993). *Maximum Entropy Models in Science and Engineering*. John Wiley & Sons, Inc., New York, USA.
- [15] Kapur, J. N., Kesavan, H. K. (1992). *Entropy optimization principles with applications*. Academic press, Inc., New York.
- [16] Noughabi, H. A. (2010). A new estimator of entropy and its application in testing normality. *Journal of Statistical Computation And Simulation* 80:1151-1162.

- [17] Paninski, L. (2003). Estimation of entropy and mutual information. *Neural Computation* 15:1191-1253.
- [18] Shannon, C.E. (1948). A mathematical theory of ommunication. *Bell System Tech. J.* 27:379-423 & 623-656.
- [19] Van Es, B. (1992). Estimating functionals related to a density by a class of statistics based on spacings. *Scandinavian Journal of Statistics* 19:61-72.
- [20] Vasicek, O. (1976). A test for normality based on sample entropy. *Journal of the Royal Statistical Society* 38:54-59.
- [21] Wiczorkowski, R. and Grzegorzewski, P. (1999). Entropy estimators improvements and comparisons. *Communications in Statistics-Simulation and Computation* 28(2):541-567.

List of Recent Technical Reports

89. Yogendra P. Chaubey, Arusharka Sen and Pranab K. Sen, *A New Smooth Density Estimator for Non-Negative Random Variables*, January 2007
90. Md. Sharif Mozumder and José Garrido, *On the Relation between the Lévy Measure and the Jump Function of a Lévy Process*, October 2007
91. Arusharka Sen and Winfried Stute, *A Bi-Variate Kaplan-Meier Estimator via an Integral Equation*, October 2007
92. C. Sangüesa, *Uniform Error Bounds in Continuous Approximations of Nonnegative Random Variables Using Laplace Transforms*, January 2008
93. Yogendra P. Chaubey, Naâmane Laib and Arusharka Sen, *A Smooth Estimator of Regression Function for Non-negative Dependent Random Variables*, March 2008
94. Alejandro Balbás, Beatriz Balbás and Antonio Heras, *Optimal Reinsurance with General Risk Functions*, March 2008
95. Alejandro Balbás, Raquel Balbás and José Garrido, *Extending Pricing Rules with General Risk Functions*, April 2008
96. Yogendra P. Chaubey and Pranab K. Sen, *On the Selection of the Smoothing Parameter in Poisson Smoothing of Histogram Estimator: Computational Aspects*, December 2008
97. Runhuan Feng and José Garrido, *Actuarial Applications of Epidemiological Models*, December 2008
98. Alejandro Balbás, Beatriz Balbás and Raquel Balbás, *CAPM and APT Like Models with Risk Measures*, June 2009
99. Jun Zhou and José Garrido, *A Loss Reserving Method Based on Generalized Linear Models*, June 2009
100. Yogendra P. Chaubey and Isha Dewan, *Smooth Estimation of Survival and Density Functions for a Stationary Associated Process Using Poisson Weights*, September 2009

101. Krishna K. Saha and Debaraj Sen, *Improved Confidence Interval for the Dispersion Parameter in Count Data Using Profile Likelihood*, September 2009
102. Araceli Reyes, *Difficulties Teaching Mathematics with the ACE Cycle*, September 2009
103. Mansi Khurana, Yogendra P. Chaubey and Shalini Chandra, *Jackknifing the Ridge Regression Estimator: A Revisit*, February 2012.
104. Yogendra P. Chaubey and Rui Zhang, *Survival Distributions with Bath-tub Shaped Hazard: A New Distribution Family*, November 2013.
105. Yogendra P. Chaubey, *Smooth Kernel Estimation of a Circular Density Function: A Connection to Orthogonal Polynomials on the Unit Circle*, January 2016.
106. Nasim Ejlali, Hamid Pezeshk, Yogendra P. Chaubey and Mehdi Sadeghi, *Parrondo's Paradox for Games with Three Players*, April 2018.
107. Yogendra P. Chaubey and Shamal C. Karmaker, *On Some Circular Distributions Induced by Inverse Stereographic Projection*, April 2018.
108. Patrice Gaillardetz and Saeb Hachem, *Risk-control Strategies*, August 2019.
109. Yogendra P. Chaubey and Nhat Linh Vu, *A Numerical Study of Entropy and Residual Entropy Estimators Based on Smooth Density Estimators for Non-negative Random Variables*, March 2020.

Copies of technical reports can be requested from:

Dr. Wei Sun
Department of Mathematics and Statistics
Concordia University
1455 de Maisonneuve Blvd. West,
Montreal, QC, H3G 1M8 CANADA